

## PATENT COOPERATION TREATY

PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

United States Patent and Trademark  
Office  
(Box PCT)  
Crystal Plaza 2  
Washington, DC 20231  
ÉTATS-UNIS D'AMÉRIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 09 June 1999 (09.06.99)	
International application No. PCT/IB98/01628	Applicant's or agent's file reference P18098WO
International filing date (day/month/year) 06 October 1998 (06.10.98)	Priority date (day/month/year) 06 October 1997 (06.10.97)
Applicant ABRIGNANI, Sergio et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

06 May 1999 (06.05.99)

☐ in a notice effecting later election filed with the International Bureau on:2. The election ☒ was☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland	Authorized officer C. Carrié
Facsimile No.: (41-22) 740.14.35	Telephone No.: (41-22) 338.83.38




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/12, 15/62, C07K 14/705, A61K 38/17, A01K 67/027, G01N 33/50, 33/576</b>		A1	(11) International Publication Number: <b>WO 99/18198</b>
		(43) International Publication Date:	15 April 1999 (15.04.99)
(21) International Application Number: PCT/IB98/01628		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 6 October 1998 (06.10.98)			
(30) Priority Data:			
9721182.5                      6 October 1997 (06.10.97)                      GB 9813560.1                      23 June 1998 (23.06.98)                      GB			
(71) Applicant (for all designated States except US): CHIRON S.P.A. [IT/IT]; Via Fiorentina, 1, I-53100 Siena (IT).			
(72) Inventors; and		<b>Published</b> With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(75) Inventors/Applicants (for US only): ABRIGNANI, Sergio [IT/IT]; Piazza Fabbrini, 5, I-53010 Vagliagli (IT). GRANDI, Guido [IT/IT]; 9A Strada, 4, I-20090 Segrate (IT).			
(74) Agent: HALLYBONE, Huw, George; Carpmals & Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB).			

(54) Title: HEPATITIS C RECEPTOR PROTEIN CD81

## (57) Abstract

The present invention relates to the use of CD81 protein and polynucleic acid in the therapy and diagnosis of hepatitis C and pharmaceutical compositions, animal models and diagnostic kits for such purposes.

1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> round screening

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## HEPATITIS C RECEPTOR PROTEIN CD81

Field of the Invention

The present invention relates to the use of CD81 protein and nucleic acid encoding this protein in the therapy and diagnosis of hepatitis C and to pharmaceutical compositions,  
5 animal models and diagnostic kits for such uses.

Brief Description of the Prior Art

All publications, manuals, patents, and patent applications cited herein are incorporated in full by reference. HCV (previously known as Non-A Non-B hepatitis - NANBV) is a positive sense RNA virus of about 10000 nucleotides with a single open reading frame  
10 encoding a polyprotein of about 3000 amino acids. Although the structure of the virus has been elucidated by recombinant DNA techniques (European patent application EP-A-0318216 and European patent application EP-A-0388232), the virus itself has not been isolated and the functions of the various viral proteins produced by proteolysis of the polyprotein have only been inferred by analogy with other similar viruses of similar  
15 genomic organisation (Choo *et al* PNAS USA (1991) 88 2451-2455).

The viral proteins are all available in recombinant form, expressed in a variety of cells and cell types, including yeast, bacteria, insect, plant and mammalian cells (Chien, D.Y. *et al* PNAS USA (1992) 89 10011-10015 and Spaete, R.R. *et al* Virology (1992) 188 819-830).

20 Two proteins, named E1 and E2 (corresponding to amino acids 192-383 and 384-750 of the HCV polyprotein respectively) have been suggested to be external proteins of the viral envelope which are responsible for the binding of virus to target cells.

HCV research is hindered very considerably by the limited host range of the virus. The only reliable animal model for HCV infection is the chimpanzee and HCV does not  
25 readily propagate in tissue culture.

In our copending International patent application PCT/IB95/00692 (WO 96/05513), we describe a method employing flow cytometry to identify cells carrying the HCV receptor. We have shown that, by labelling cells with recombinant E2 envelope protein,

it is possible to sort cells using flow cytometry, isolating those cells capable of specific binding to the E2 and therefore potentially carrying an HCV receptor.

In our copending International patent application PCT/IB96/00943 (WO 97/09349), we have identified a protein capable of binding to the E2 envelope protein of HCV.

- 5 We have now succeeded with some difficulty in cloning the DNA encoding the HCV receptor and have discovered, surprisingly that the DNA encodes a cellular protein known as CD81. We are not aware of any association in the literature between CD81 and the HCV. CD81 was first identified by monoclonal antibodies as the target of an antiproliferative antibody (TAPA-1) which inhibited *in vitro* cellular proliferation.
- 10 Armed with this new information and given the sequence knowledge of CD81 in the public databases it is now possible to design and produce an armoury of therapeutic and diagnostic reagents against HCV.

#### Summary of the Invention

- According to the present invention, there is provided a CD81 protein, or functional equivalent thereof, for use in the therapy or diagnosis of hepatitis C (HCV). According to a further aspect of the present invention there is provided a compound that binds specifically to the CD81 protein for use in the therapy or diagnosis of HCV.
- 15

- The term "CD81 protein, or a functional equivalent thereof" as used herein means the human CD81 protein as defined by the protein sequence listed in the SWISSPROT database (Accession No. P18582) or the EMBL/GENBANK database (Accession No. M33690 ) or a functional equivalent thereof. A functional equivalent of CD81 is a compound which is capable of binding to HCV, preferably to the E2 protein of HCV. Preferably, the functional equivalent is a peptide or protein. The term "functional equivalent" includes an analogue of CD81, a fragment of CD81, and CD81 mutants and muteins.
- 20
- 25

One region of the human CD81 protein that is shown herein to be involved in binding to the E2 protein of HCV is the "EC2" region comprising amino acids 113-201 of the full length human sequence shown in Figure 1. The invention encompasses proteins and protein fragments containing this region of human CD81, or containing functional

equivalents of this region, such as, for example, the Chimpanzee sequence identified in Figure 1. Preferably, the functional equivalent is at least 80% homologous to the human CD81 sequence across the EC2 region of the protein, preferably at least 90% homologous as assessed by any conventional analysis algorithm such as for example, the Pileup sequence analysis software (Program Manual for the Wisconsin Package, 1996).

The term "a functionally equivalent fragment" as used herein also means any fragment or assembly of fragments of the complete protein that binds to HCV, preferably that binds to the E2 protein of HCV. The complete protein may be truncated at one or both ends or domains may be removed internally provided that the protein retains the defined function. For example, one or more regions of the protein responsible for membrane binding (TM1 to TM4 in Figure 1) may be removed to render the protein soluble when produced by a recombinant process. The fragment of choice comprises the extracellular loop 2 (EC2 in Figure 1) of the CD81 protein (amino acids 113-201).

If proteinaceous, functionally equivalent fragments or analogues may belong to the same protein family as the human CD81 protein identified herein. By "protein family" is meant a group of proteins that share a common function and exhibit common sequence homology. By sequence homology is meant that the protein sequences are related by divergence from a common ancestor, such as is the case between the human and the chimpanzee. Chimpanzee CD81 is thus an example of a functionally equivalent protein that binds to HCV.

Preferably, the homology between functionally equivalent protein sequences is at least 25% across the whole of amino acid sequence of the complete protein or of the complete EC2 fragment (amino acids 113-201). More preferably, the homology is at least 50%, even more preferably 75% across the whole of amino acid sequence of the protein or protein fragment. Most preferably, homology is greater than 80% across the whole of the sequence.

The term "a functionally equivalent analogue" is used to describe those compounds that possess an analogous function to an activity of the CD81 protein and may, for example comprise a peptide, cyclic peptide, polypeptide, antibody or antibody fragment. These compounds may be proteins, or may be synthetic agents designed so as to mimic certain

structures or epitopes on the inhibitor protein. Preferably, the compound is an antibody or antibody fragment.

The term "functionally equivalent analogue" also includes any analogue of CD81 obtained by altering the amino acid sequence, for example by one or more amino acid deletions, substitutions or additions such that the protein analogue retains the ability to bind to HCV, preferably the E2 protein of HCV. Amino acid substitutions may be made, for example, by point mutation of the DNA encoding the amino acid sequence.

The functional equivalent of CD81 may be an analogue of a fragment of CD81. The CD81 or functional equivalent may be chemically modified, provided it retains its ability to bind to HCV, preferably the E2 protein of HCV.

It is envisaged that such molecules will be extremely useful in preventative therapy of HCV infection, because these molecules will bind specifically to the virus and will thus prevent internalisation of the virus into cells. As used herein, "binding specifically" means that the functionally equivalent analogue has high affinity for the E2 protein of the HCV virus and does not bind to any other protein with similar high affinity. Specific binding may be measured by a number of techniques such as Western blotting, FACS analysis, or immunoprecipitation. Preferably, the functionally equivalent analogue binds to the E2 protein with an affinity of at least  $10^{-8}$ , preferably at least  $10^{-9}$  and most preferably greater than  $10^{-10}$ .

According to a further embodiment of the invention there is provided a compound that binds to CD81, such as a monoclonal or polyclonal antibody to CD81, for use in the diagnosis or therapy of HCV. Preferably the compound binds specifically to CD81 with an affinity of at least  $10^{-8}$ , preferably at least  $10^{-9}$  and most preferably greater than  $10^{-10}$ . Such compounds may be used to prevent the virus binding to patient cells and being internalised.

The CD81 molecule is present on a number of different cell types. Ideally, the compound that binds to CD81 therefore only interacts with CD81 in the presence of HCV, so that the usual function of CD81 is not compromised on healthy cells. Antibodies and suitable methods of screening for such antibodies are described in co-pending applications EP 96928648.3 and EP 95927918.3.

The CD81 protein, or functional equivalent thereof may be produced by any suitable means, as will be apparent to those of skill in the art. In order to produce sufficient amounts of CD81 protein, or functional equivalents thereof for use in accordance with the present invention, expression may conveniently be achieved by culturing under  
5 appropriate conditions recombinant host cells containing the CD81 protein, or functional equivalent thereof.

Systems for cloning and expression of a polypeptide in a variety of different host cells are well known.

Two preferred methods of construction of carrier proteins according to the invention are  
10 direct chemical synthesis and by production of recombinant protein. Preferably, the CD81 protein is produced by recombinant means, by expression from an encoding nucleic acid molecule. Recombinant expression has the advantage that the production of the protein is inexpensive, safe, facile and does not involve the use of toxic compounds that may require subsequent removal.

15 When expressed in recombinant form, the CD81 protein or functional equivalent thereof is preferably generated by expression from an encoding nucleic acid in a host cell. Any host cell may be used, depending upon the individual requirements of a particular system. Suitable host cells include bacteria, mammalian cells, plant cells, yeast and baculovirus systems. Mammalian cell lines available in the art for expression of a  
20 heterologous polypeptide include Chinese hamster ovary cells, HeLa cells, baby hamster kidney cells and many others. Preferably, bacterial hosts are used for the production of recombinant protein, due to the ease with which bacteria may be manipulated and grown. A common, preferred bacterial host is *E. coli*.

Preferably, if produced recombinantly, the CD81 protein or functional equivalent is  
25 expressed from a plasmid that contains a synthetic nucleic acid insert. The insertion site in the expression plasmid into which the nucleic acid encoding the CD81 protein or functional equivalent is cloned may allow linkage of the protein to a tag, such as the "flag" peptide or polyhistidine. This arrangement facilitates the subsequent purification of recombinant protein.



According to a further aspect of the present invention, there is also provided a nucleic acid molecule encoding the CD81 protein or functional equivalent thereof, for use in the therapy or diagnosis of HCV infection. Preferably, the nucleic acid encodes human CD81 protein. As will be apparent to one of skill in the art, such a nucleic acid molecule  
5 will be designed using the genetic code so as to encode the protein or peptide that is desired. A nucleic acid molecule according to this aspect of the present invention may comprise DNA, RNA or cDNA and may additionally comprise nucleotide analogues in the coding sequence. Preferably, the nucleic acid molecule will comprise DNA.

Nucleotide sequences included within the scope of this embodiment of the invention are  
10 those hybridising to nucleic acid encoding the CD81 protein under standard conditions. As used herein, standard conditions includes both non-stringent standard hybridisation conditions (6 x SSC/50% formamide at room temperature) with washing under conditions of low stringency (2 x SSC/50% formamide at room temperature, or 2 x SSC, 42°C) or at standard conditions of higher stringency, e.g. 2 x SSC, 65°C (where SSC = 0.15M NaCl,  
15 0.015M sodium citrate, pH 7.2). Preferably the term standard conditions refers to conditions of high stringency.

Preferably, such nucleic acid molecules will retain the ability to hybridise specifically to nucleic acid encoding CD81 or a fragment thereof and will include nucleic acid sequences with 40% homology across the whole of the human CD81 gene sequence as  
20 defined by the Pileup command of the GCG Program manual for the Wisconsin Package (version 9, 1996). More preferably, the homology is at least 65% across the whole of the gene sequence. Most preferably, homology is greater than 70% across the whole of the gene sequence.

Nucleic acid encoding the CD81 protein or functional equivalent may be cloned under  
25 the control of an inducible promoter, so allowing precise regulation of protein expression. Suitable inducible systems will be well known to those of skill in the art.

Suitable vectors for the expression of the CD81 protein or functional equivalent may be selected from commercial sources or constructed in order to suit a particular expression system. Such vectors will contain appropriate regulatory sequences, such as promoter  
30 sequences, terminator sequences, polyadenylation sequences, enhancer sequences and marker genes. Vectors may be plasmids, or viral-based. For further details see

Molecular Cloning: a laboratory manual (Sambrook *et al.*, 1989). Many known techniques and protocols for the manipulation of nucleic acids and analysis of proteins are described in detail in "Short protocols in molecular biology", second addition, Ausubel *et al.* (John Wiley & Sons 1992).

- 5 Methods for the isolation and purification of recombinant proteins will be well known to those of skill in the art and are summarised, for example in Sambrook *et al* (1989). Particularly in bacteria such as *E. coli*, the recombinant protein will form inclusion bodies within the bacterial cell, thus facilitating its preparation. If produced in inclusion bodies, the carrier protein may need to be refolded to its natural conformation.
- 10 Additionally, in order to tailor precisely the exact properties of the CD81 protein or functional equivalent thereof, the skilled artisan will appreciate that changes may be made at the nucleotide level from known CD81 sequences, by addition, substitution, deletion or insertion of one or more nucleotides. Site-directed mutagenesis (SDM) is the method of preference used to generate mutated proteins according to the present invention.
- 15 There are many techniques of SDM now known to the person of skill in the art, including oligonucleotide-directed mutagenesis using PCR as set out, for example by Sambrook *et al.*, (1989) or using commercially available kits.

Suitable vectors can be chosen or constructed, containing appropriate regulatory sequences, including promoter sequences, terminator sequences, polyadenylation sequences, enhancer sequences, marker genes and other sequences as appropriate.

20 Vectors may be plasmids, viral e.g. 'phage, or phagemid, as appropriate. For further details see, for example, *Molecular Cloning: a Laboratory Manual*: 2nd edition, Sambrook *et al.*, 1989, Cold Spring Harbor Laboratory Press. Many known techniques and protocols for manipulation of nucleic acid, for example in preparation of nucleic acid constructs, mutagenesis, sequencing, introduction of DNA into cells and gene

25 expression, and analysis of proteins, are described in detail in *Short Protocols in Molecular Biology*, Second Edition, Ausubel *et al.* eds., John Wiley & Sons, 1992. The disclosures of Sambrook *et al.* and Ausubel *et al.* are incorporated herein by reference.

- 30 According to a further aspect of the invention, there is provided a method for treating an infection of HCV comprising administering to a patient a therapeutically effective

amount of CD81 protein, or a functional equivalent thereof effective to reduce the infectivity of the virus.

Since the infection mechanism of HCV appears to depend, in part, upon the availability of a cell surface receptor, making available a soluble form of the CD81 protein, or a functional equivalent thereof will act as an antagonist of binding of HCV to the cellular receptor thus reducing or preventing the infection process and thereby treating the disease.

A suitable soluble form of the CD81 protein, or a functional equivalent thereof might comprise, for example, a truncated form of the protein from which one or more of the transmembrane domain or domains TM1-TM4 have been removed either by a protein cleavage step or, by design, in a chemical or recombinant DNA synthesis. The preferred soluble form of the protein comprises the EC2 domain (residues 113-201 as identified in Figure 1). The EC1 domain may act to increase the affinity or specificity of the protein for HCV.

Alternatively, a hybrid particle comprising at least one particle-forming protein, such as hepatitis B surface antigen or a particle-forming fragment thereof, in combination with the CD81 protein or functional equivalent thereof could be used as an antagonist of binding of HCV to the cellular receptor.

According to a still further aspect of the invention, there is provided a method for treating an infection of HCV comprising administering to a patient a therapeutically effective amount of a compound that specifically binds to CD81 protein, such as a monoclonal antibody directed to CD81. The rationale behind this therapeutic strategy is that the binding of the cell surface receptor to another compound will prevent the binding of HCV to the receptor, so preventing the infection process and thereby treating the disease.

According to a further aspect of the invention, there is provided a pharmaceutical composition comprising a CD81 protein or functional equivalent thereof, optionally as a pharmaceutically acceptable salt, in combination with a pharmaceutically acceptable carrier. According to a still further aspect of the present invention there is provided a pharmaceutical composition comprising a compound that binds specifically to the CD81

protein, optionally as a pharmaceutically acceptable salt, in combination with a pharmaceutically acceptable carrier.

The pharmaceutical composition may be in any appropriate form for administration including oral, parenteral, transdermal and transcutaneous compositions. The  
5 composition may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated.

A process is also provided for making the pharmaceutical composition, in which a protein of the present invention is brought into association with a pharmaceutically acceptable carrier.

10 According to a further aspect of the invention, there is provided a CD81 protein or functional equivalent thereof, or a compound that binds specifically to the CD81 protein for use as a pharmaceutical.

According to a further aspect of the invention, there is provided the use of a CD81 protein or functional equivalent thereof or compound that binds specifically to the  
15 CD81 protein in the manufacture of a medicament for the treatment of an HCV infection.

The ability of a CD81 protein or functional equivalent thereof to bind to HCV permits the use of the protein as a diagnostic for HCV infection, for example in an ELISA (Enzyme linked immunosorbent assay) or RIA (Radioimmunoassay).

20 A soluble form of the protein could, for example, be used in an ELISA form of assay to measure neutralising antibodies in serum. More preferably, antibodies to CD81 will be suitable for use in this context, since these molecules will be anti-idiotypic antibodies for HCV itself.

According to a further aspect of the invention, there is provided an assay for HCV  
25 antibodies in a serum sample comprising the step of allowing competitive binding between antibodies in the sample and a known amount of an HCV protein for binding to a CD81 protein or functional equivalent thereof and measuring the amount of the known HCV protein bound.

Preferably, the CD81 protein or functional equivalent thereof is immobilised on a solid support and the HCV protein, which may suitably be E2 HCV envelope protein, optionally recombinant E2 protein, is labelled. The label may be a radioactive label, a peptide, an epitope, an enzyme, or any other bioactive compound. Preferably the label  
5 comprises an enzyme.

In an assay of this form, competitive binding between antibodies and the HCV protein for binding to the CD81 protein or functional equivalent thereof results in the bound HCV protein being a measure of antibodies in the serum sample, most particularly, HCV neutralising antibodies in the serum sample.

10 A significant advantage of the assay is that direct measurement is made of neutralising of binding antibodies (i.e. those antibodies which interfere with binding of HCV envelope protein to the cellular receptor). Such an assay, particularly in the form of an ELISA test has considerable applications in the clinical environment and in routine blood screening.

15 Also, since the assay measures neutralising of binding antibody titre, the assay forms a ready measure of putative vaccine efficacy, neutralising of binding antibody titre being correlated with host protection.

In a further aspect of the invention, there is provided a diagnostic kit comprising the CD81 protein or functional equivalent thereof. Preferably the kit also contains at least  
20 one labelled HCV protein, optionally enzyme labelled. The kit will also contain other components necessary for the analysis of the presence of HCV or anti-HCV antibodies in serum. Such components will be readily apparent to those of skill in the art.

The CD81 protein or functional equivalent thereof may be used to screen for chemical compounds mimicking the HCV surface structure responsible for binding to the HCV  
25 receptor.

According to a further aspect of the invention, there is provided a method for screening chemical compounds for ability to bind to the region of HCV responsible for binding to a host cell, comprising measuring the binding of a chemical compound to be screened to

a CD81 protein or functional equivalent thereof . The host cell may be any mammalian cell, preferably a human host cell.

This aspect of the invention encompasses the products of the screening process whether alone, in the form of a pharmaceutically acceptable salt, in combination with one or  
5 more other active compounds and/or in combination with one or more pharmaceutically acceptable carriers. Processes for making a pharmaceutical composition are also provided in which a chemical compound identified by the process of the invention is brought into association with a pharmaceutically acceptable carrier.

The chemical compound may be an organic chemical and may contain amino acids or  
10 amino acid analogues. Preferably however the chemical compound is a peptide, polypeptide or a polypeptide which has been chemically modified to alter its specific properties, such as the affinity of binding to the CD81 protein or functional equivalent thereof or its stability *in vivo*.

According to a further aspect of the invention, there is provided a nucleic acid encoding  
15 CD81 protein or functional equivalent thereof for use in diagnosis or therapy of HCV. The nucleic acid may encode any part of the CD81 protein, or functional equivalent thereof. Preferably, the nucleic acid encodes a portion of CD81 that binds to HCV E2. According to a still further aspect of the present invention, there is provided a nucleic acid encoding a peptide or polypeptide compound that binds specifically to CD81.

20 Changes to the nucleic acid may be made at the nucleotide level by addition, substitution, deletion or insertion of one or more nucleotides, which changes may or may not be reflected at the amino acid level, dependent on the degeneracy of the genetic code.

The nucleic acid may be included in a vector, optionally an expression vector permitting  
25 expression of the nucleic acid in a suitable host to produce CD81 protein or functional equivalent thereof.

The identification of the DNA encoding the HCV receptor, namely CD81, makes available the full power of molecular biology for the molecular analysis of HCV and in particular its infectious mechanism, offering for the first time the possibility of

designing methods of treating the virus. PCR methods may be used to identify cells carrying the receptor and DNA molecules may be designed to act as polymerase chain reaction (PCR) primers in this connection. Although CD81 is widespread and is associated with normal human function, the present invention includes antisense  
5 molecules inhibiting CD81 production for use in the treatment of HCV and in the manufacture of a medicament for the treatment of HCV infection.

The identification of polymorphisms in the CD81 protein may be found to be associated with susceptibility to HCV infection or likely prognosis. Accordingly, the identification of the gene encoding the HCV receptor allows the evaluation of polymorphisms present  
10 throughout the human population.

According to a further aspect of the invention, there is provided an antibody to CD81 protein or functional equivalent thereof for use in the treatment of an HCV infection and in the manufacture of a medicament for the treatment of an HCV infection. The antibody is preferably a monoclonal antibody. Such an antibody can be used to  
15 temporarily block the CD81 receptor preventing infection from HCV, for example, immediately after an accidental infection with HCV-infected blood.

At present, the only available animal model of HCV infection is the chimpanzee, which is a protected species. Experiments on such animals pose a number of difficulties which together result in a very considerable expense (a one year experiment with one  
20 chimpanzee can cost \$70,000). Compared to this, a mouse model would be far more acceptable. Unfortunately, as described below, the HCV receptor, whilst ubiquitous in humans and found in chimpanzees, is absent in other mammals. A transgenic mammal, for example a mouse, carrying the HCV receptor on the cell surface, perhaps expressed in greater or lesser amounts than normally found, would be of great benefit to HCV  
25 research and the development of vaccines. Expression of mutant CD81 proteins on the surface of cells would also be a useful research tool.

According to a further aspect of the invention, there is provided a transgenic non-human animal, suitably a mammal such as a mouse, carrying a transgene encoding a CD81 protein or functional equivalent thereof.

The transgenic animal of the invention may carry one or more other transgenes to assist in maintaining an HCV infection.

There is also provided a process for producing a transgenic animal comprising the step of introducing a DNA encoding a CD81 protein or functional equivalent thereof into the  
5 embryo of a non-human mammal, preferably a mouse. Preferably the CD81 protein or functional equivalent thereof is a human CD81 protein.

According to a further aspect of the present invention, there is provided a CD81 protein or a functional equivalent thereof for use as a protective immunogen in the control of HCV.

#### 10 Brief Description of the Drawings

Figure 1 is a sequence alignment showing the homology between human, chimpanzee, green monkey, hamster, rat and mouse CD81 gene sequences.

Figure 1A is a schematic description of primary, secondary and tertiary rounds of screening.

15 Figure 1B is a schematic description of the final round of screening.

Figure 2 is a FACS scan analysis of E2 bound cells.

Figure 3 shows the dose-dependent inhibition of anti-CD81 binding to B cells by recombinant E2. The data are expressed as % inhibition of mean fluorescence intensity.

Figure 4 is an immunoblot showing the recognition of the membrane protein fraction  
20 immunoprecipitated by anti-CD81 antibody. Lane 2: recombinant E2 precipitated with chimpanzee antiserum to E2; lane 3, recombinant E2 precipitated with chimpanzee pre-immune serum lane 4: 20µg of anti-CD81 mAb (clone JS81 Pharmingen) precipitated with goat anti-mouse IgG, lane 5: control, (20 µg of an irrelevant monoclonal antibody, anti-human CD9, ATCC) precipitated with goat anti-mouse IgG  
25 linked to protein A sepharose. Lane 1: positive control, membrane protein preparation.



Figure 5 shows the nucleotide and deduced amino acid sequences of the EC2 fragment cloned in pThio-His C and the upstream plasmid sequence coding for the carboxyl terminus of thioredoxin and for the enterokinase cleavage site.

Figure 6 shows the appearance of a protein band of the expected molecular mass for thioredoxin-EC2 in the extract from the induced sample.

Figure 7 is a Coomassie Blue stained gel showing the purification of thioredoxin-EC2.

Figure 8 represents the nucleotide and deduced amino acid sequence of the EC2-His<sub>6</sub> fragment cloned into pGEX-KG as well as the upstream plasmid sequence coding for the carboxyl terminus of GST, the thrombin cleavage site and a small glycine spacer.

Figure 9 represents an SDS-PAGE of total proteins of the TOP10 *E. coli* clone which express GST-EC2-(His)<sub>6</sub>.

Figure 10 is a Coomassie-stained SDS-PAGE showing thrombin cleavage of GST-EC2-(His)<sub>6</sub> after purification of the protein on a glutathione sepharose column.

Figure 11 shows the dose-dependent inhibition of E2 binding to hepatocarcinoma cells by recombinant molecule expressing the major extracellular loop (EC2) of human CD81.

Figure 12 shows binding of HCV to CD81.

Figures 13-17 show the construction of nucleic acid vectors for use in the generation of mice transgenic for the human CD81 gene.

20

### Detailed Description of the Invention

#### **Example 1. Recombinant E2, cell lines, vector DNA, and antibodies used in the present study.**

The recombinant E2 used in this screening was produced in CHO cells (E2-CHO) (WO 97/09349). E2-CHO binds to the human T cell lymphoma cell line Molt-4. A subline of Molt-4 (termed A2A6), was identified by expanding individual Molt-4 cell colonies

and testing for the amount of E2-CHO that bound to the cell surface. The A2A6 subline was found to bind more E2-CHO molecule on its surface than its parental line and was therefore chosen for the source of RNA, expecting that this subline may have a higher representation of the transcript encoding the E2 binding molecule. These cells were  
5 chosen using an assay whereby human B and T lymphoma cells and hepatocarcinoma cell lines were incubated with recombinant E2 expressed in mammalian cells (CHO) as described by D. Rosa *et al.*, *Proc. Natl. Acad. Sci. USA* **93**, 1759 (1996) and stained with biotin-labelled anti-E2 antibodies as described by Rosa *et al.*, (1996). Cells with the highest E2 binding ability were sorted using a FacsVantage (Becton Dickinson) and  
10 subcloned by limiting dilution. Growing clones were screened for E2 binding at the Facs and clones with the highest Mean Fluorescence Intensity were further expanded.

WOP is a NIH3T3-derived cell which expresses polyoma T antigen (L. Dailey and C. Basilico, *J. Virol.* **54**, 739 (1985). In this cell line, plasmids containing the polyoma replication origin can be amplified episomally. Recombinant DNA constructed with  
15 pCDM8 (Invitrogen) can be recovered from selected transfectants, which contains the polyoma replication origin and is designed for the manipulation of expression libraries in eukaryotic cells.

A mouse monoclonal anti-E2 antibody (291A2) was used for detection of E2-CHO bound on the cell surface of transfectants. This antibody was obtained as follows:  
20 BALB-c mice were immunised three times with recombinant E2 (10µg) in complete Freund's adjuvant. Cell fusions between spleen cells and non-producing myeloma cells were made according to standard techniques. The supernatant from fusions was then screened for binding to E2 bound to Molt-4 cells, so as to identify monoclonal antibodies that bound to an exposed site on the E2 molecule. The most suitable antibody  
25 identified in this fashion was termed 291A2.

### **Example 2: Construction of cDNA library**

Total RNA was extracted from the A2A6 cell line according to the method described by Chomczynsky and Sacchi (Chomczynsky, P. and Sacchi, N. (1987) *Anal. Biochem.* **162**: 156-159). Poly(A)+ was enriched twice using oligo(dT) cellulose. Starting from 2µg of  
30 this RNA as a template, the double strand complementary DNA was synthesized using a Superscript II cDNA synthesis kit (Life Technologies) in the presence of oligo(dT)

(100ng) and random hexamer primers (100ng). The cDNA was blunt-ended with T4 DNA polymerase, and was ligated with a *Bst*XI linker, which allows the insertion of the fragment into the same restriction site in the polylinker region of the expression vector pCDM8. The linker-ligated cDNA was phenol-extracted and ethanol precipitated using ammonium sulphate to remove free mononucleotides. followed by Sephacryl 500 chromatography (Lifetechnologies) to size-fractionate the cDNA. The purified cDNA fragment over 500bp were pooled and ligated with *Bst*XI - digested pCDM8 at a molecular ratio of approximately 1:1. This final ligation reaction was used from transformation of *E. coli* MC1061/P3 by electroporation using Gene-Pulser (BIORAD).  
10 A total of  $2 \times 10^6$  cfu was amplified and pooled in liquid bacterial culture as a cDNA library.

### Example 3: Library screening

The screening procedure was based largely on the method described by Campbell *et al.* (Campbell, I. G., Jones, T.A., Foulkes, W. D. and Trowsdale, J. Cancer Res. 51: 5329-5338, 1991). Enrichment was carried out using magnetic beads (the first to the third round) (Figure 1A) and panning techniques (the fourth round). (Figure 1B).

#### 3.1 The first round of screening

A total of 375µg of amplified DNA, which represents  $2 \times 10^6$  of independent cDNA clones, was prepared. In each transfection, 25µg of DNA was mixed with  $10^7$  WOP cells using the Gene-Pulser electroporator (BIORAD) under the conditions of 300V/500µF. Fifteen sets of transfections were performed. After transfection, cells were incubated at 37°C for 2 days and then the cells were detached by trypsinization and washed with PBS supplemented with 5% FCS and 0.5mM EDTA twice by centrifugation at 360 x g for 10min at 4°C. The cell pellet was resuspended in PBS supplemented with 5% FCS and 0.5mM EDTA ( $10^7$  cells/ml) and then E2-CHO was added to the cell suspension at a concentration of 10µg/ml. The cells were incubated on ice for 60 min. After washing twice with PBS supplemented with 5% FCS and 0.5mM EDTA, the cell suspension was incubated with 291A2 antibody on ice for 30 min. After washing twice with PBS supplemented with 5% FCS and 0.5mM EDTA, 10µl of Dynabeads (DYNAL) coupled with goat anti-mouse IG was added to the cell suspension. The mixture was gently agitated using a Coulter Mixer (Coulter) for 60

min at 4°C. Bound cells were separated using Magnetic Particle Concentrator (DYNAL) from non-binders, according to the manufacturer's instructions, thus enriching E2-binding transfectants. Plasmid DNA was recovered from the bound transfectated cells using the protocol described by Campbell *et al.* (Campbell, I. G., Jones, T.A., Foulkes, W. D. and Trowsdale, J. Cancer Res. 51: 5329-5338, 1991). *E. coli* MC1061/P3 was transformed with this plasmid by electroporation. This DNA pool is referred to as the first enriched pool (1°EP).

### 3.2 The second round of screening

A total of 150µg of amplified DNA derived from 1°EP was prepared and 6 sets of the transfection were performed and transfectants were enriched using the same condition as in the first screening. This DNA pool is referred to as 2°EP.

### 3.3 The third round of screening

A total of 25µg of amplified DNA derived from 2°EP was prepared and one set of the transfection was performed. Transfectants were enriched using the same condition as in the first screening. During this separation step, transfectants formed aggregates, which might be caused by expression of irrelevant adhesion molecules. This could decrease the efficiency of enrichment because these aggregates contained magnetic beads non-specifically. To circumvent this potential problem, transfectants after the second separation by Magnetic Particle Concentrator were diluted and plated on Terasaki plates. Approximately 100 of single cells identified under microscope were pooled and plasmid DNA was extracted from them. The DNA pool prepared from this step is referred to as 3°EP.

### 3.4 The fourth round of screening

291A1 monoclonal antibody was incubated in a Petri dish (90mm) at a concentration of 10µg/ml overnight at 4°C.

A total of 25µg of amplified DNA derived from 3°EP was prepared and one set of transfections was performed. The transfectated cells were incubated with E2-CHO as described above, and placed onto the 291A2-coated plates for 60 min at 4°C. After rinsing with a large excess of PBS supplemented with 5% FCS and 0.5mM EDTA

twice, the bound cells were directly treated with the lysing solution and plasmids were extracted as described as before. This DNA pool is referred to as 4°EP.

### 3.4 Identification of cDNA encoding a molecule binding to the recombinant E2

DNA was isolated from single colonies derived from 4°EP. A single transfection was performed for each plasmid preparation using the same conditions as used for the previous screening steps. E2-binding of the transformants was detected using a phycoerythrin-conjugated monoclonal Fab fragment of goat anti-mouse Ig instead of the antibody-coupled Dynabeads. Transfectants of 3°EP and 4°EP were also analyzed in the same way. The E2-bound cells were detected on FACScan (Becton Dickinson) and analyzed with LYSIS II program (Becton Dickinson) (Figure 2). E2-CHO binds increasingly as the purification step advances. A single clone P3 showed strong E2-binding.

#### **Example 4: DNA sequencing determination and analysis.**

P3 contains a insert of approximately 1 kb. The DNA sequence of the insert of the cDNA clone which confers E2-binding to WOP upon transfection was determined by an automated sequencing system using the T7 primer, whose sequence is located adjacent the cloning site of pCDM8. The sequence was screened through the GenBank databases using the GCG programs on a UNIX computer. This analysis revealed that the 5' part of P3 insert is identical to human CD81 (TAPA-1). Restriction analysis of P3 using three enzymes (*Bst*XI), *Hinc*II and *Nco*I) also agreed with the restriction map of human CD81 cDNA.

#### **Example 5: Binding of CD81 to recombinant E2.**

Anti-CD81 antibodies were used to assess the interaction between E2 and CD81. EBV-B cells were incubated with increasing concentrations of recombinant E2 for 1 hour at 4°C and then stained with an anti-CD81 monoclonal antibody (clone JS-81, Pharmingen). As shown in Figure 3, recombinant E2 was found to competitively inhibit the binding of anti-CD81 antibodies to EBV transformed B-cell lines (EBV-B cells). The data are expressed as % inhibition of mean fluorescence intensity (Rosa *et al.*, 1996).

In addition, E2 reacts in Western blot with anti-CD81 precipitated material (Figure 4). This Figure shows E2 recognition of membrane protein fraction immunoprecipitated by anti-CD81 antibody. Approximately 300 µg of membrane protein extract prepared from the A2A6 cell line were solubilised in 8mM CHAPS in PBS pH 7.4, incubated with 10 µg recombinant E2 (lanes 2 and 3), with 20µg of anti-CD81 mAb (clone JS81; Pharmingen) (lane 4), or as control, with 20 µg of an irrelevant monoclonal antibody (anti-human CD9, ATCC) (lane 5) for 2 hours at 4°C, and finally precipitated with chimpanzee antiserum to E2 (lane 2), chimpanzee pre-immune serum (lane 3), or goat anti-mouse IgG (lanes 4 and 5) bound to protein A sepharose (CL-4B, Pharmacia). The pellet was dissolved in Laemmli buffer and subjected to SDS-PAGE under non-reducing conditions. After electroblotting, the PVDF membrane (Millipore) was incubated overnight with 1µg/ml of recombinant E2 at room temperature, and for 2 hours with 291A2 anti-E2 monoclonal antibody. E2 binding to immunoprecipitated proteins was detected with an anti-mouse IgG peroxidase-conjugated polyclonal antibody (Amersham). As a positive control membrane proteins also were loaded on the gels (lane 1). The mobility of molecular weight standards is indicated on the left in kilodaltons.

CD81 is also expressed on fresh lymphocytes and hepatocytes as demonstrated by immunohistochemical staining with biotin-labelled-E2 or anti-CD81 (data not shown).

To assess whether CD81 could mediate the internalisation of ligands, we exploited the fact that CD81 forms a complex with CD19 and CD21 on the surface of B lymphocytes (D. T. Fearon and R. H. Carter, 1995, *Annu. Rev. Immunol.* 13, 127). B cells were incubated with E2 at 37°C for different times, after which CD19 or CD21 levels on the cell surface were measured by immunofluorescence. Incubation of B cells with E2 resulted in down-regulation of both CD19 and CD21 (data not shown). It thus seems as if CD81 is able to mediate the internalisation of both these ligands.

**Example 6: The major extracellular loop of CD81 binds recombinant E2 and viral particles.**

To map the CD81 domain that binds E2 protein our efforts were focused on the EC2 hydrophilic extracellular loop of the protein. This fragment was expressed in *E. coli* as a Thioredoxin-EC2 fusion protein that has an enterokinase site between thioredoxin and

EC2, and as a GST-EC2 fusion protein which has a thrombin site between GST and EC2 and a hexa-histidine tag added to the carboxyl-terminus of the protein. We show that both proteins are expressed and are able to bind HCV E2. In competition experiments we also show that the purified fusion proteins and the EC2-His fragment  
 5 excised from GST-thrombin-EC2-(His)<sub>6</sub> are able to inhibit the binding of E2 on the surface of CD81 expressing cells.

#### 6.1 Cloning of EC2 in pThio-His.

Figure 5 shows the nucleotide and the deduced amino acid sequences of the EC2 fragment cloned in pThio-His C and the upstream plasmid sequence coding for the  
 10 carboxyl terminus of thioredoxin and for the enterokinase cleavage site. As shown, EC2 is fused in frame with thioredoxin through the enterokinase site, which can be exploited to remove thioredoxin from the fusion protein.

The fragment coding for EC2 was PCR-amplified from the plasmid pCDM8/P3 using the following oligodeoxynucleotides:

15 Forward BL EC2  
 5'GGCGGGGGTGGATCCGGGGGTGGAGGCTCGAGCTTTGTCAACAAGGACC3  
 XhoI Phe Val Asn Lys Asp  
 Reverse BL EC2  
 5'CCCCAAGCTT TCA CAG CTT CCC GGA GAA GAG GTC ATC G3'  
 20 HindIII Stop Leu Lys Gly Ser Phe Leu Asp Asp

Using standard cloning techniques (Sambrook *et al.*, 1989) the PCR product was double-digested with *XhoI* and *HindIII*, ligated to pThio-His C (Invitrogen) digested with the same restriction enzymes, and transformed into Top10 *E. coli* cells. After selection of the transformants by restriction enzyme analysis and DNA sequencing of  
 25 the plasmids, a correct construct coding for the expected thioredoxinenterokinase site-EC2 fusion protein was identified. Glycerol batches of selected clones were stored to -80°C.

Total protein extracts of the thioredoxin-EC2 expressing clone before and after IPTG addition, were subjected to SDS-PAGE to analyse protein expression. Figure 6 clearly shows the appearance of a protein band of the expected molecular mass (23.4 kDa) in the extract from the induced sample. The figure also shows the reactivity of the fusion protein with E2. The TOP10 *E. coli* clone containing the pThio-hisC-EC2 plasmid and a TOP10 clone containing the pThio-HisC plasmid devoid of insert were induced, soluble protein extracts were prepared from both clones and subjected to Far Western Blot with E2 protein. For this blot, protein samples were brought to 1x loading sample buffer (LSB) (5% w/v SDS, 10% v/v glycerol, 62.5 mM Tris-HCl, 0.05 % Bromophenol Blue) using a 3x LSB solution. The samples were run onto a 15% polyacrylamide gel and transferred to a PVDF membrane (Immobilon-P, Millipore). The membrane was incubated for 30 min in blocking solution (PBS, 10% w/v non-fat dried milk, 0.05% v/v Tween 20). Following an incubation of 15 hours at 4°C with blocking solution containing 1 µg/ml of CHO-E2, the membranes were incubated for 2 hours with the 291A2 anti-E2 monoclonal antibody diluted 1:250, and for 1 hour with a peroxidated goat antimouse Ig antibody (Sigma) diluted 1:2000. Three washing steps between all incubation steps were performed using blocking solution, which was also used to dilute the antibodies. After a final wash with PBS the membranes were incubated for 1 min with luminol (ECL, Amersham) and exposed on Hyper-film (Amersham).

As can be seen from these Figures, a band corresponding to the molecular weight of Thioredoxin-EC2 was visible in the lane where the soluble proteins from the pThio-HisC-EC2 were loaded. Such a band was absent in the lane where the soluble proteins of the pThio-HisC clone were loaded.

## 6.2 Purification of Thioredoxin-EC2

For the purification of thioredoxin-EC2 the following procedure was developed:

- 1) osmotic shock of the cells, 2) protein precipitation with 30% saturation ammonium sulphate, and 3) IMAC. After osmotic shock about 50% of the fusion protein was released from the cells together with contaminant proteins. The ammonium sulphate precipitation resulted in a pellet which contained thioredoxin-EC2 devoid of the bulk of contaminant proteins. IMAC of the resuspended precipitate resulted in a fusion protein



which was about 85% pure as assessed by SDS-PAGE. With this procedure we purified 5 mg thioredoxin-EC2 from a litre of culture. This procedure is set out in detail below.

The *E. coli* clone expressing Thioredoxin-EC2 was inoculated in 500 ml LB medium containing 100 µg/ml ampicillin. At  $OD_{600} = 0.5$ , 0.5 mM IPTG was added to the culture and growth was continued at 37°C for additional 3.5 hours. The culture was then centrifuged at 4000 x g for 10 min at 4°C, the cell pellet was resuspended with 50 ml ice cold hypertonic solution (20 mM Tris-HCl, 2.5 mM EDTA, 20 % sucrose, pH 8) and left on ice for 10 min. The resuspended cells were centrifuged again as above and the pellet was resuspended in hypotonic buffer (20 mM Tris-HCl, 2.5 mM EDTA, pH 8) to osmotically shock the cells. After 20 min at 0°C the suspension was centrifuged at 12.000 x g for 10 min at 4°C, the supernatant was brought to 30%  $NH_4(SO_4)_2$  using a room temperature saturated solution of the salt. The suspension was incubated overnight at 4°C and then centrifuged at 10.000 x g for 10 min. The pellet was resuspended using 15 ml of 20 mM Phosphate buffer, 500 mM NaCl, pH 6, clarified by centrifugation, and loaded on a 2 ml column of Nickel activated Chelating Sepharose Fast Flow (Pharmacia) equilibrated in the same buffer.

After adsorption, the column was washed with 10 ml of the equilibrium buffer (flow rate 0.5 ml/min), and then the Thioredoxin-EC2 was eluted using a 30 ml gradient 0-50 mM Imidazole in 20 mM Phosphate buffer, 500 mM NaCl, pH 6 followed by an isocratic elution with 10 ml of 400 mM imidazole. 2.4 ml fractions were collected. The fractions containing the recombinant protein were pooled, dialysed against PBS, and stored to -20 °C. Proteins were analysed by means of SDS-PAGE and protein content was assayed by the Bradford method using BSA as a protein standard.

Purified Thioredoxin-EC2 is shown in Figure 7.

### 6.3 Cloning of EC2-(His)<sub>6</sub> in pGEX-KG

Figure 8 represents the nucleotide and deduced amino acid sequence of the EC2-(His)<sub>6</sub> fragment cloned in pGEX-KG as well as the upstream plasmid sequence coding for the carboxyl terminus of GST, the thrombin cleavage site, and a small glycine spacer. As shown, EC2 is fused in frame with GST through the thrombin site, which can be exploited to remove GST from the fusion protein. The glycine-rich spacer, located

between thrombin site and EC2, facilitates the cleavage of the fusion protein by thrombin (Guan, K.L., and Dixon, J.E. (1991) *Anal. Biochem.* **192**, 262-267).

The fragment coding for EC2 was PCR-amplified from the plasmid pCDM8/P3 using the following oligodeoxynucleotides:

5            EC2 Forward            EC2

5' CAAAAGGAATTCTA TTT GTC AAC AAG GAC CAG ATC GCC AAG3'

*EcoRI*            Phe Val Asn Lys Asp Gln Ile Ala Lys

Reverse BLH

His tag

EC2

5'CCCCAAGCTTTCAATGATG ATG ATG ATG ATG CAG CTT CCC GGA  
10 GAAG3'

*HindIII* Stop His His His His His His Leu Lys Gly Ser  
Phe

The PCR product was digested with *XhoI* and *HindIII*, ligated to pGEX-KG (Guan, K. L., and Dixon, J. E. (1991) *Anal. Biochem.* **192**, 262-267) digested with the same  
15 restriction enzymes, and transformed into TOP10 *E. coli* cells. After selection of the transformants by restriction enzyme analysis and nucleotide sequencing of the plasmids, a plasmid having the expected size of the insert was found to have also the correct EC2-(His)<sub>6</sub> sequence in frame with the upstream thrombin and GST coding sequence. The plasmid prepared from the selected TOP10 clone was then transformed into BL21 cells.  
20 Glycerol batches of selected clones were stored to -80°C.

Figure 9 represents an SDS-PAGE of total proteins of the TOP10 *E. coli* clone which expresses GST-EC2-(His)<sub>6</sub>. This analysis clearly shows that in the extract of the induced sample a protein band with the expected molecular mass (39kDa) was present. The corresponding Far Western Blot clearly shows the E2 specifically reacts with the  
25 fusion protein.

#### 6.4 Purification of GST-EC2-(His)<sub>6</sub>

The GST-EC2-(His)<sub>6</sub> fusion protein was purified on a glutathione sepharose column and digested with thrombin (Figure 10). After digestion, the EC2-(His)<sub>6</sub> moiety was further purified by two additional chromatographic steps consisting of a glutathione sepharose  
5 column to remove the GST fragment and IMAC chromatography. This procedure is detailed below.

A single colony of an *E. coli* clone expressing the GST-EC2 fusion protein was inoculated in 10 ml LB, 100 µg/ml Amp and cells were grown overnight at 37 °C. The culture was then inoculated in 500 ml of medium and when OD<sub>600</sub> = 0.5 was reached  
10 0.5 mM IPTG was added. After 3.5 hours the cells were harvested by centrifugation, resuspended with 9 ml PBS and disrupted with two passages at 18.000 psi using a French Press (SLM Aminco). The lysate was centrifuged at 30.000xg and the supernatant was loaded on a column of 1 ml of Glutathione Sepharose 4B (Pharmacia) equilibrated in PBS.

15 The column was washed with 10 ml PBS, and eluted with 4 ml of 50 mM Tris-HCl, 10mM reduced glutathione, pH 8. The eluted proteins were dialysed against PBS and stored to -20°C.

#### 6.5 Digestion of GST-EC2-(His)<sub>6</sub> with thrombin and purification of EC2-(His)<sub>6</sub>

9.6 mg of protein recovered from the glutathione sepharose column were digested with  
20 22 units of thrombin (Pharmacia) for 8 hours at room temperature, then the enzyme was inactivated using 0.13 mM PMSF (Sigma). The reaction mixture was then dialysed against PBS and loaded into 0.5 ml of GST-sepharose column equilibrated in PBS. The column was washed with 1 ml of PBS. The flow-through and the wash were pooled and loaded into 0.250 ml of Nickel-activated chelating sepharose column. EC2-(His)<sub>6</sub> was  
25 recovered from the column eluting with 1 ml of 20 mM phosphate buffer, 500 mM NaCl, 400 mM imidazole, pH 7.8. A dialysis was then performed against PBS.

#### **Example 7: Binding of CD81 fragment to virus**

The proteins containing the human, but not the mouse EC2 loop of CD81, bound to E2 in western blot (data not shown) and inhibited binding of E2 to human cells (Figure 11).

The chimeric proteins were coated on polystyrene beads and incubated with an infectious plasma containing known amounts of viral RNA molecules. After washing, the bead-associated virus was assessed by quantitative RT-PCR for the amount of bound HCV RNA. This experiment was performed as set out below.

- 5 Polystyrene beads (1/4 inch diameter) (Pierce) were coated overnight with purified EC2 recombinant protein in citrate buffer pH4 at room temperature. After saturation for one hour with 2% BSA in 50mM TrisCl pH 8, 1mM EDTA, 100mM NaCl (TEN) buffer, each bead was incubated at 37°C for 2 hours in 200µl TEN-diluted infectious chimp plasma containing  $5 \times 10^5$  HCV RNA molecules.
- 10 For inhibition experiments, the EC2-coated polystyrene beads were incubated with 10µg/ml of purified monoclonal antibodies for one hour at room temperature before incubation with the virus. Each bead was washed 5 times with 15ml TEN buffer in an automated washer (Abbot) and viral RNA was extracted using the Viral Extraction Kit (Qiagen). RNA (8 ml) was reverse-transcribed at 42°C for 90 minutes in 20 ml Buffer A
- 15 (Perkin Elmer Taq Man) containing 100pmol of the HCV antisense primer CGGTTCCGCAGACCACTATG, 40 U RNAsin (Promega), 5 nmol dNTPs, 110 nmol MgCl<sub>2</sub>, 10U M-MuRT (Boheringer). cDNA (20 ml) was amplified using a Perkin-Elmer ABI 7700 Sequence Detection System (45 cycles) in 50 ml Buffer A containing 100 pmol of the HCV sense primer TCTTCACGCAGAAAGCGTCTA, 5 pmol of the
- 20 fluorescent detection probe 5'(FAM)TGAGTGTCGTGCAGCCTCCAGGA(TAMRA) (kindly provided by David Slade, Pharmacia and Upjohn), 15 nmol dNTPs, MgCl<sub>2</sub> and 1.25U Taq Gold (Perkin-Elmer, Foster City, CA). All reactions were quantified using HCV (genotype 1a) infected plasma (bDNA titer of 30 mEq/ml) to generate a standard curve. Sequence Detector Software from Perkin-Elmer has been previously described
- 25 (U. E. Gibson, C. A. Heid and P. M. Williams, *Genome Res.* 6, 995 (1996)).

As shown in Figure 12, the molecules containing the human CD81 extracellular loop bound HCV in a concentration-dependent fashion, and pre-incubation of the chimeric proteins with anti-CD81 antibodies inhibited virus binding. Furthermore, serum from chimpanzees which were protected from homologous challenge by vaccination with

30 recombinant E1/E2 envelope heterodimer (Q.-L. Choo *et al. Proc. Natl. Acad. Sci. USA*

91. 1294 (1994)) completely inhibited HCV binding to bead-coated-CD81, while serum from vaccinated and non-protected animals did not (data not shown).

These data demonstrate that expression of human CD81, and in particular its major extracellular loop are sufficient for binding not only E2 but also HCV particles. Given  
5 the wide distribution of CD81 (S. Levy, S. C. Todd and H. T. Maecker, *Annu. Rev. Immunol.* **16**, 89 (1998), these results imply that HCV binds and may be internalised by a variety of cells other than hepatocytes. Indeed, HCV RNA has been found in T and B lymphocytes and monocytes (K. Blight, R. R. Lesniewski, J. T. LaBrooy and E. J. Gowans, *Hepatology* **20**, 553 (1994); P. Bouffard *et al.*, *J. Infect. Dis.* **166**, 1276 (1992);  
10 Zignego *et al.*, *J. Hepatol.* **15**, 382 (1992)). Whether virus binding is followed by entry and infection in all cell types is not clear because of the lack of an efficient HCV culture system in vitro. It may well be that CD81 is an HCV attachment receptor and that additional factors are required for viral fusion or infectivity.

CD81 participates in different molecular complexes on different cell types, a fact that  
15 may influence its capacity to serve as a receptor for HCV infection or to deliver regulatory signals to target cells. For instance, it associates with integrins on epithelial and hematopoietic cells (F. Berditchevski, M. Zutter and M. E. Hemler, *Mol. Biol. Cell* **7**, 193 (1996); B. A. Mannion, F. Berditchevski, S.-K. Kraeft, L. B. Chen and M. E. Hemler, *J. Immunol.* **157**, 2039 (1996)), whereas it is part of a signaling complex  
20 containing CD21, CD19 and Leu13 on B cells (L. E. Bradbury, G. S. Kansas, S. Levy, R. L. Evans and T. F. Tedder, *J. Immunol.* **149**, 2841 (1991)). This complex has been shown to facilitate antigen specific stimulation by lowering the activation threshold of B cells (D. T. Fearon and R. H. Carter, *Annu. Rev. Immunol.* **13**, 127 (1995)). It is worth  
25 noting that HCV appears to use a molecule that is part of the same complex containing the EBV receptor (CD21) (N. R. Cooper, M. D. Moore and G. R. Nemerow, *Annu. Rev. Immunol.* **6**, 85 (1988)), and the ability of EBV to activate and immortalise B lymphocytes is well documented.

#### **Example 8: Construction of transgenes**

The following constructs were designed and made in order to generate mice transgenic  
30 for human CD81.

1. Addition of splicing and polyadenylation signals of rabbit beta-globin gene to the human CD81 cDNA fragment.

The human CD81 cDNA fragment from the pCDM8/P3 clone was transferred into a pBluescript KS II(+) vector (Stratagene) and was then inserted into the pSPP plasmid (derived from BMGSC expression vector, a kind gift from Dr. Karasuyama, Basel Institute for Immunology) between two fragments, one containing the second intron and the other containing the polyadenylation signal of the rabbit beta-globin gene (position 902-1547 and 1543-2081, respectively, GenBank accession No. M12603) (pSR1P in Figure 11). The resulting recombinant DNA fragment was excised from the pBluescript KSII(+) vector (Stratagene) by *SalI* (at 5' end) and *BamHI* (at 3' end).

2. Creation of a transgene for ubiquitous expression of human CD81

The *SalI*-*BamHI* fragment of the pSR1P insert was inserted into the compatible restriction sites of pCAGmcs, a modified plasmid of pCAGGS (a kind gift from Dr. J. Miyazaki at Osaka University, Japan, under restricted permission), which contains chicken beta-actin promoter and human cytomegalovirus enhancer (Niwa, H. *et al.*, Gene 108, p193 (1991). (pCAGSR1Pp in Figure 12). The 3.8 kb *EcoRI*-*BamHI* fragment was submitted to zygote injection.

3. Creation of a transgene for liver-specific expression of human CD81

The *SalI* site of pSR1P was converted to a *BamHI* site by *BamHI* linker ligation after blunt-end formation with Klenow fragment of *E. coli* DNA polymerase I. This *BamHI* fragment was inserted into the *BamHI* site of the ALB *e/p* plasmid, carrying the mouse albumin promoter and enhancer (Pinkert, C.A. *et al.*, Genes Dev. 1, p268 (1987) (received from Dr. F. Chisari, Scripps Research Institute, La Jolla, San Diego). (pAlbSR1P in Figure 13) The 4.5 kb *NotI*-*EcoRV* fragment was submitted to zygote injection.

4. Creation of a transgene for B lymphocyte-specific expression of human CD81

700 bp *BamHI* fragment of the mouse immunoglobulin heavy chain enhancer (a kind gift from Dr. A. Kudo, Basel Institute for Immunology) and 2.3 kb *XbaI*-*SacI* fragment

of the mouse kappa light chain promoter was subcloned into a pBluescript KSII(+) vector. The *SacI* site was converted to a *HindIII* site by *HindIII* linker ligation described above. The *BamHI* site of pCAGSR1P was first converted to *NotI* site. Then the promoter region of the modified pCAGSR1P construct was removed by *EcoRI*-  
5 *HindIII* restriction digestion and replaced with the immunoglobulin promoter-enhancer fragment. (pEhKpSR1P in Figure 15) The 5.2 kb *EcoRI*-*BamHI* fragment was submitted to zygote injection.

Together, our data indicate that CD81 is an attachment receptor for HCV and may  
10 provide new insight into the mechanisms of HCV infection pathogenesis. Since CD81 associates with an activation complex on the surface of B cells, the present finding may explain the pathogenesis of HCV associated cryoglobulinemia, even if there is no viral replication in B cells. Moreover, the identification of the interaction between HCV and CD81 may help in mapping conserved neutralising epitopes on the virus envelope  
15 which should be important to develop effective vaccines and to provide a decoy receptor for viral neutralisation.

## CLAIMS

- 1 A CD81 protein, or a functional equivalent thereof for use in the therapy or diagnosis of HCV.
- 2 A protein comprising the human CD81 sequence listed in the SWISSPROT  
5 database (Accession No. P18582) or the EMBL/GENBANK database (Accession No. M33690) or a functional equivalent thereof for use in the therapy or diagnosis of HCV.
- 3 A protein comprising an amino acid sequence with at least 80% homology to the human CD81 sequence listed in the SWISSPROT database (Accession No.  
10 P18582) or the EMBL/GENBANK database (Accession No. M33690), homology being defined using the Pileup sequence analysis software package (Wisconsin, 1996), for use in the therapy or diagnosis of HCV.
- 4 A protein comprising amino acids 113-201 of the human CD81 sequence listed in the SWISSPROT database (Accession No. P18582) or the EMBL/GENBANK  
15 database (Accession No. M33690), or a functional equivalent thereof.
- 5 A protein according to claim 4, for use in the therapy or diagnosis of HCV.
- 6 A compound that binds specifically to a CD81 protein, for use in the therapy or diagnosis of HCV.
- 7 A method for treating an infection of HCV comprising administering to a  
20 patient a therapeutically effective amount of a CD81 protein, or a functional equivalent thereof or administering a compound that binds specifically to the CD81 protein, to reduce the infectivity of the virus.
- 8 A pharmaceutical composition comprising a CD81 protein, or a functional equivalent thereof, or a compound that binds specifically to a CD81 protein,  
25 optionally as a pharmaceutically acceptable salt, in combination with a pharmaceutically acceptable carrier.



- 9 A pharmaceutical composition comprising a protein according to claim 4 in combination with a pharmaceutically acceptable carrier.
- 10 A pharmaceutical composition according to either of claims 8 or 9 for use in the therapy or diagnosis of HCV.
- 5 11 A process for preparing a pharmaceutical composition as defined in claim 8 or 9, in which a CD81 protein, or a functional equivalent thereof, or a protein according to claim 4 or a compound that binds specifically to a CD81 protein is brought into association with a pharmaceutically acceptable carrier.
- 10 12 Use of a CD81 protein, a functional equivalent thereof or a compound that binds specifically to a CD81 protein in the manufacture of a medicament for the treatment or diagnosis of an HCV infection.
- 13 Use of a protein according to claim 4 in the manufacture of a medicament for the treatment or diagnosis of an HCV infection.
- 15 14 An assay for HCV antibodies present in a serum sample comprising the step of allowing competitive binding between antibodies in the sample, a known amount of HCV protein and a known amount of a CD81 protein, or a functional equivalent thereof and measuring the amount of the known HCV protein that binds to the CD81 protein.
- 20 15 An assay for HCV in a serum sample comprising the step of allowing competitive binding between antibodies in the sample and a known amount of a CD81 protein, or a functional equivalent thereof and measuring the amount of the known CD81 protein bound.
- 16 A diagnostic kit comprising a CD81 protein, or a functional equivalent thereof, optionally labeled.
- 25 17 A diagnostic kit according to claim 16 wherein the label comprises a radioactive label, a peptide, an epitope, an enzyme, or other bioactive compound,
- 18 A method for screening chemical compounds for ability to bind to the region of HCV responsible for binding to a host cell, comprising measuring the binding of a

chemical compound to be screened to a CD81 protein, or a functional equivalent thereof.

19 A transgenic non-human mammal, carrying a transgene encoding CD81 protein, or a functional equivalent thereof.

5 20 A process for producing a transgenic animal comprising the step of introducing a DNA encoding a CD81 protein into the embryo of a non-human mammal, preferably a mouse.

21 A nucleic acid molecule which encodes a CD81 protein, or a functional equivalent thereof for use in the treatment or diagnosis of HCV.

10 22 A nucleic acid molecule which hybridises to a nucleic acid molecule as defined in claim 21 under standard conditions.

23 A nucleic acid molecule which hybridises to a nucleic acid molecule as defined in claim 21 under conditions of high stringency (2 x SSC, 65°C).

15 24 The nucleic acid molecule according to any of claims 21-23 which comprises DNA.

25 A CD81 protein or a functional equivalent thereof for use as a protective immunogen in the control of HCV.

26 A fusion protein comprising a CD81 protein or functional equivalent thereof for use in the treatment or diagnosis of HCV.

---CY1---||---TM1---||---EC1---||---TM2---||---CY2---||---TM3---

MGVEGCTKCIKYILLFVENFVPHLAGGVILGVALWLRHDPOTTNLLYLELGDKRPAPNTFFVVGIIYLLAVGAVNMFVGLGCYGAJQESQCLLGTPTCLVI<sup>100</sup>

**Human  
Chimpanzee  
Green Monkey  
Hamster  
Rat  
Mouse**

SECRET

	260
101	101
102	102
103	103
104	104
105	105
106	106
107	107
108	108
109	109
110	110
111	111
112	112
113	113
114	114
115	115
116	116
117	117
118	118
119	119
120	120
121	121
122	122
123	123
124	124
125	125
126	126
127	127
128	128
129	129
130	130
131	131
132	132
133	133
134	134
135	135
136	136
137	137
138	138
139	139
140	140
141	141
142	142
143	143
144	144
145	145
146	146
147	147
148	148
149	149
150	150
151	151
152	152
153	153
154	154
155	155
156	156
157	157
158	158
159	159
160	160
161	161
162	162
163	163
164	164
165	165
166	166
167	167
168	168
169	169
170	170
171	171
172	172
173	173
174	174
175	175
176	176
177	177
178	178
179	179
180	180
181	181
182	182
183	183
184	184
185	185
186	186
187	187
188	188
189	189
190	190
191	191
192	192
193	193
194	194
195	195
196	196
197	197
198	198
199	199
200	200

Human  
Chimpanzee  
Green Monkey  
Hamster  
Rat  
Mouse


-----T44-----|-----CY3-----|

101 278  
K L Y L G I A A I V V A V I H I F E M I L S H T L C C G I R N S S V  
.....  
.....  
.....  
.....


Human  
Chimpanzee  
Green Monkey  
Hamster  
Rat  
Mouse

**FIG. 1**

2/17

**FIG. 1A****1<sup>o</sup>, 2<sup>o</sup>, 3<sup>o</sup> round screening**

# FIG. 1B <sup>3/17</sup> 4<sup>c</sup> round screening






FIG. 2B




FIG. 2A

5/17

FIG. 4





FIG. 3






**FIG. 5**




FIG. 6




8/17

**FIG. 7****FIG. 10**


**FIG. 8**

9/17

**FIG. 9**



11/17

**FIG. 11**

12/17






FIG. 12

13/17


**FIG. 13**

14/17


**FIG. 14**




15/17

**FIG. 15**

16/17

**FIG. 16**

17/17

**FIG. 17**

# INTERNATIONAL SEARCH REPORT

national Application No

PCT/IB 98/01628

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC 6 C12N15/12 C12N15/62 C07K14/705 A61K38/17 A01K67/027  
G01N33/50 G01N33/576

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K A01K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	OREN R ET AL: "TAPA-1, THE TARGET OF AN ANTIPROLIFERATIVE ANTIBODY, DEFINES A NEW FAMILY OF TRANSMEMBRANE PROTEINS" MOLECULAR AND CELLULAR BIOLOGY, vol. 10, no. 8, August 1990, pages 4007-4015, XP000749300 see the whole document ---	1-5, 21-25
X	ANDRIA M L ET AL: "GENOMIC ORGANIZATION AND CHROMOSOMAL LOCALIZATION OF THE TAPA-1 GENE" JOURNAL OF IMMUNOLOGY, vol. 147, no. 3, 1 August 1991, pages 1030-1036, XP002064247 see the whole document --- -/--	1-5, 21-25

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

8 February 1999

Date of mailing of the international search report

18/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hornig, H

# INTERNATIONAL SEARCH REPORT

national Application No

PCT/IB 98/01628

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 09349 A (BIOCINE SPA ; ABRIGNANI SERGIO (IT)) 13 March 1997 cited in the application see the whole document ----	6
X	EP 0 318 216 A (CHIRON CORP) 31 May 1989 cited in the application see the whole document ----	6
X	EP 0 388 232 A (CHIRON CORP) 19 September 1990 cited in the application see the whole document ----	6
A	ROSA D ET AL: "A QUANTITATIVE TEST TO ESTIMATE NEUTRALIZING ANTIBODIES TO THE HEPATITIS C VIRUS: CYTOFLUORIMETRIC ASSESSMENT OF ENVELOPE GLYCOPROTEIN 2 BINDING TO TARGET CELLS" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 93, no. 3, March 1996, pages 1759-1763, XP000615446 cited in the application see the whole document ----	1-26
A	WO 96 05513 A (BIOCINE SPA ; ABRIGNANI SERGIO (IT)) 22 February 1996 cited in the application see the whole document ----	1-26
T	P. PILERI ET AL.: "Binding of hepatitis C virus to CD81" SCIENCE, vol. 282, 30 October 1998, pages 938-941, XP002092549 AAAS, WASHINGTON, DC, US see the whole document -----	1-26

# INTERNATIONAL SEARCH REPORT

international application No.

PCT/IB 98/01628

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 7 and 20  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/IB 98/01628

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9709349	A	13-03-1997	AU 6835196 A CA 2227301 A EP 0852586 A	27-03-1997 13-03-1997 15-07-1998
EP 0318216	A	31-05-1989	AU 2796789 A CN 1035679 A CN 1073719 A DD 287104 A DD 298524 A DD 298525 A DD 298526 A DD 298527 A DE 3886363 D DE 3886363 T DK 353789 A ES 2012739 T FI 893447 A FI 981380 A GB 2212511 A, B GR 90300069 T HK 38293 A HR 940493 A HU 9500427 A IE 62868 B IN 171237 A IN 171238 A IN 171239 A IN 171240 A IN 169067 A JP 2810032 B JP 10108674 A JP 10290696 A JP 10290697 A JP 2662358 B JP 6225797 A JP 2810022 B JP 9184844 A JP 9173079 A JP 2809956 B JP 6128290 A JP 2662350 B JP 6128291 A JP 2532805 B JP 6128292 A JP 2662351 B JP 6128293 A JP 2500880 T LV 10726 A LV 10726 B NO 955101 A NO 955102 A PT 89041 A, B SI 8812138 A US 5714596 A	14-06-1989 20-09-1989 30-06-1993 14-02-1991 27-02-1992 27-02-1992 27-02-1992 27-02-1992 27-01-1994 19-05-1994 18-07-1989 01-12-1994 17-07-1989 15-06-1998 26-07-1989 31-07-1991 30-04-1993 31-08-1997 28-12-1995 08-03-1995 22-08-1992 22-08-1992 22-08-1992 22-08-1992 31-08-1991 15-10-1998 28-04-1998 04-11-1998 04-11-1998 08-10-1997 16-08-1994 15-10-1998 15-07-1997 08-07-1997 15-10-1998 10-05-1994 08-10-1997 10-05-1994 11-09-1996 10-05-1994 08-10-1997 10-05-1994 29-03-1990 20-06-1995 20-12-1995 15-12-1995 15-12-1995 01-12-1988 31-08-1997 03-02-1998
EP 0388232	A	19-09-1990	AU 4750493 A AU 666767 B AU 666576 B	20-01-1994 22-02-1996 15-02-1996

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 98/01628

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0388232 A		AU 640920 B	09-09-1993
		AU 5278390 A	22-10-1990
		CA 2012482 A	17-09-1990
		DD 297446 A	09-01-1992
		FI 981381 A	15-06-1998
		GR 91300016 T	15-11-1991
		HU 9500746 A	29-01-1996
		JP 2693908 B	24-12-1997
		JP 7101986 A	18-04-1995
		JP 2809987 B	15-10-1998
		JP 7145194 A	06-06-1995
		JP 10309197 A	24-11-1998
		JP 10295389 A	10-11-1998
		JP 9215497 A	19-08-1997
		JP 2656995 B	24-09-1997
		JP 4504715 T	20-08-1992
		NO 303941 B	28-09-1998
		NO 960741 A	23-02-1996
		NO 960742 A	23-02-1996
		PT 93480 A, B	07-11-1990
		US 5714596 A	03-02-1998
		WO 9011089 A	04-10-1990
		US 5679342 A	21-10-1997
		US 5350671 A	27-09-1994
		US 5698390 A	16-12-1997
		US 5712088 A	27-01-1998
		US 5863719 A	26-01-1999
		US 5683864 A	04-11-1997
WO 9605513 A	22-02-1996	AU 3189495 A	07-03-1996
		CA 2174212 A	22-02-1996
		EP 0723665 A	31-07-1996
		JP 9504377 T	28-04-1997



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : <b>C12N 15/12, 15/62, C07K 14/705, A61K 38/17, A01K 67/027, G01N 33/50, 33/576</b>		<b>A1</b>	(11) International Publication Number: <b>WO 99/18198</b>
			(43) International Publication Date: 15 April 1999 (15.04.99)
(21) International Application Number: <b>PCT/IB98/01628</b>		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 6 October 1998 (06.10.98)			
(30) Priority Data: 9721182.5 6 October 1997 (06.10.97) GB 9813560.1 23 June 1998 (23.06.98) GB			
(71) Applicant (for all designated States except US): <b>CHIRON S.P.A. [IT/IT]; Via Fiorentina, 1, I-53100 Siena (IT).</b>			
(72) Inventors; and (75) Inventors/Applicants (for US only): <b>ABRIGNANI, Sergio [IT/IT]; Piazza Fabbrini, 5, I-53010 Vagliagli (IT). GRANDI, Guido [IT/IT]; 9A Strada, 4, I-20090 Segrate (IT).</b>			
(74) Agent: <b>HALLYBONE, Huw, George; Carpmaels &amp; Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB).</b>			

Published

With international search report.


Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: **HEPATITIS C RECEPTOR PROTEIN CD81**

(57) Abstract

The present invention relates to the use of CD81 protein and polynucleic acid in the therapy and diagnosis of hepatitis C and pharmaceutical compositions, animal models and diagnostic kits for such purposes.

1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> round screening



# INTERNATIONAL SEARCH REPORT

national Application No

PCT/IB 98/01628

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC 6 C12N15/12 C12N15/62 C07K14/705 A61K38/17 A01K67/027  
G01N33/50 G01N33/576

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K A01K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	OREN R ET AL: "TAPA-1, THE TARGET OF AN ANTIPROLIFERATIVE ANTIBODY, DEFINES A NEW FAMILY OF TRANSMEMBRANE PROTEINS" MOLECULAR AND CELLULAR BIOLOGY, vol. 10, no. 8, August 1990, pages 4007-4015, XP000749300 see the whole document	1-5, 21-25
X	ANDRIA M L ET AL: "GENOMIC ORGANIZATION AND CHROMOSOMAL LOCALIZATION OF THE TAPA-1 GENE" JOURNAL OF IMMUNOLOGY, vol. 147, no. 3, 1 August 1991, pages 1030-1036, XP002064247 see the whole document	1-5, 21-25
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"8" document member of the same patent family

Date of the actual completion of the international search

8 February 1999

Date of mailing of the international search report

18/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Hornig, H

# INTERNATIONAL SEARCH REPORT

national Application No

PCT/IB 98/01628

**C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 09349 A (BIOCINE SPA ; ABRIGNANI SERGIO (IT)) 13 March 1997 cited in the application see the whole document	6
X	EP 0 318 216 A (CHIRON CORP) 31 May 1989 cited in the application see the whole document	6
X	EP 0 388 232 A (CHIRON CORP) 19 September 1990 cited in the application see the whole document	6
A	ROSA D ET AL: "A QUANTITATIVE TEST TO ESTIMATE NEUTRALIZING ANTIBODIES TO THE HEPATITIS C VIRUS: CYTOFLUORIMETRIC ASSESSMENT OF ENVELOPE GLYCOPROTEIN 2 BINDING TO TARGET CELLS" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 93, no. 3, March 1996, pages 1759-1763, XP000615446 cited in the application see the whole document	1-26
A	WO 96 05513 A (BIOCINE SPA ; ABRIGNANI SERGIO (IT)) 22 February 1996 cited in the application see the whole document	1-26
T	P. PILERI ET AL.: "Binding of hepatitis C virus to CD81" SCIENCE, vol. 282, 30 October 1998, pages 938-941, XP002092549 AAAS, WASHINGTON, DC, US see the whole document	1-26

# INTERNATIONAL SEARCH REPORT

international application No.

PCT/IB 98/01628

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 7 and 20  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/IB 98/01628

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9709349	A	13-03-1997	AU 6835196 A CA 2227301 A EP 0852586 A	27-03-1997 13-03-1997 15-07-1998
EP 0318216	A	31-05-1989	AU 2796789 A CN 1035679 A CN 1073719 A DD 287104 A DD 298524 A DD 298525 A DD 298526 A DD 298527 A DE 3886363 D DE 3886363 T DK 353789 A ES 2012739 T FI 893447 A FI 981380 A GB 2212511 A, B GR 90300069 T HK 38293 A HR 940493 A HU 9500427 A IE 62868 B IN 171237 A IN 171238 A IN 171239 A IN 171240 A IN 169067 A JP 2810032 B JP 10108674 A JP 10290696 A JP 10290697 A JP 2662358 B JP 6225797 A JP 2810022 B JP 9184844 A JP 9173079 A JP 2809956 B JP 6128290 A JP 2662350 B JP 6128291 A JP 2532805 B JP 6128292 A JP 2662351 B JP 6128293 A JP 2500880 T LV 10726 A LV 10726 B NO 955101 A NO 955102 A PT 89041 A, B SI 8812138 A US 5714596 A	14-06-1989 20-09-1989 30-06-1993 14-02-1991 27-02-1992 27-02-1992 27-02-1992 27-02-1992 27-01-1994 19-05-1994 18-07-1989 01-12-1994 17-07-1989 15-06-1998 26-07-1989 31-07-1991 30-04-1993 31-08-1997 28-12-1995 08-03-1995 22-08-1992 22-08-1992 22-08-1992 22-08-1992 31-08-1991 15-10-1998 28-04-1998 04-11-1998 04-11-1998 08-10-1997 16-08-1994 15-10-1998 15-07-1997 08-07-1997 15-10-1998 10-05-1994 08-10-1997 10-05-1994 11-09-1996 10-05-1994 08-10-1997 10-05-1994 29-03-1990 20-06-1995 20-12-1995 15-12-1995 15-12-1995 01-12-1988 31-08-1997 03-02-1998
EP 0388232	A	19-09-1990	AU 4750493 A AU 666767 B AU 666576 B	20-01-1994 22-02-1996 15-02-1996

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 98/01628

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0388232 A		AU 640920 B	09-09-1993
		AU 5278390 A	22-10-1990
		CA 2012482 A	17-09-1990
		DD 297446 A	09-01-1992
		FI 981381 A	15-06-1998
		GR 91300016 T	15-11-1991
		HU 9500746 A	29-01-1996
		JP 2693908 B	24-12-1997
		JP 7101986 A	18-04-1995
		JP 2809987 B	15-10-1998
		JP 7145194 A	06-06-1995
		JP 10309197 A	24-11-1998
		JP 10295389 A	10-11-1998
		JP 9215497 A	19-08-1997
		JP 2656995 B	24-09-1997
		JP 4504715 T	20-08-1992
		NO 303941 B	28-09-1998
		NO 960741 A	23-02-1996
		NO 960742 A	23-02-1996
		PT 93480 A, B	07-11-1990
		US 5714596 A	03-02-1998
		WO 9011089 A	04-10-1990
		US 5679342 A	21-10-1997
		US 5350671 A	27-09-1994
		US 5698390 A	16-12-1997
		US 5712088 A	27-01-1998
		US 5863719 A	26-01-1999
		US 5683864 A	04-11-1997
WO 9605513 A	22-02-1996	AU 3189495 A	07-03-1996
		CA 2174212 A	22-02-1996
		EP 0723665 A	31-07-1996
		JP 9504377 T	28-04-1997

## PCT

## INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference <b>P18098W0</b>	<b>FOR FURTHER ACTION</b> see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. <b>PCT/IB 98/01628</b>	International filing date (day/month/year) <b>06/10/1998</b>	(Earliest) Priority Date (day/month/year) <b>06/10/1997</b>
Applicant <b>CHIRON S.P.A. et al.</b>		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

1. ☒ Certain claims were found unsearchable (see Box I).
2. ☐ Unity of invention is lacking (see Box II).
3. ☒ The international application contains disclosure of a **nucleotide and/or amino acid sequence listing** and the international search was carried out on the basis of the sequence listing
  - ☐ filed with the international application.
  - ☒ furnished by the applicant separately from the international application,
    - ☐ but not accompanied by a statement to the effect that it did not include matter going beyond the disclosure in the international application as filed.
  - ☐ Transcribed by this Authority
4. With regard to the title,
  - ☐ the text is approved as submitted by the applicant
  - ☒ the text has been established by this Authority to read as follows:  
**HEPATITIS C RECEPTOR PROTEIN CD81**
5. With regard to the abstract,
  - ☒ the text is approved as submitted by the applicant
  - ☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this International Search Report, submit comments to this Authority.
6. The figure of the drawings to be published with the abstract is:
  - Figure No. 1A
    - ☐ as suggested by the applicant.
    - ☒ because the applicant failed to suggest a figure.
    - ☐ because this figure better characterizes the invention.
  - ☐ None of the figures.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB 98/01628

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
**Remark: Although claims 7 and 20**  
**are directed to a method of treatment of the human/animal**  
**body, the search has been carried out and based on the alleged**  
**effects of the compound/composition.**
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 98/01628

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C12N15/62 C07K14/705 A61K38/17 A01K67/027  
G01N33/50 G01N33/576

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K A01K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	OREN R ET AL: "TAPA-1, THE TARGET OF AN ANTIPROLIFERATIVE ANTIBODY, DEFINES A NEW FAMILY OF TRANSMEMBRANE PROTEINS" MOLECULAR AND CELLULAR BIOLOGY, vol. 10, no. 8, August 1990, pages 4007-4015, XP000749300 see the whole document ---	1-5, 21-25
X	ANDRIA M L ET AL: "GENOMIC ORGANIZATION AND CHROMOSOMAL LOCALIZATION OF THE TAPA-1 GENE" JOURNAL OF IMMUNOLOGY, vol. 147, no. 3, 1 August 1991, pages 1030-1036, XP002064247 see the whole document --- -/--	1-5, 21-25



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## ° Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

8 February 1999

Date of mailing of the international search report

18/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hornig, H

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 98/01628

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 09349 A (BIOCINE SPA ;ABRIGNANI SERGIO (IT)) 13 March 1997 cited in the application see the whole document ---	6
X	EP 0 318 216 A (CHIRON CORP) 31 May 1989 cited in the application see the whole document ---	6
X	EP 0 388 232 A (CHIRON CORP) 19 September 1990 cited in the application see the whole document ---	6
A	ROSA D ET AL: "A QUANTITATIVE TEST TO ESTIMATE NEUTRALIZING ANTIBODIES TO THE HEPATITIS C VIRUS: CYTOFLUORIMETRIC ASSESSMENT OF ENVELOPE GLYCOPROTEIN 2 BINDING TO TARGET CELLS" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 93, no. 3, March 1996, pages 1759-1763, XP000615446 cited in the application see the whole document ---	1-26
A	WO 96 05513 A (BIOCINE SPA ;ABRIGNANI SERGIO (IT)) 22 February 1996 cited in the application see the whole document ---	1-26
T	P. PILERI ET AL.: "Binding of hepatitis C virus to CD81" SCIENCE, vol. 282, 30 October 1998, pages 938-941, XP002092549 AAAS, WASHINGTON, DC, US see the whole document -----	1-26

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 98/01628

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9709349	A	13-03-1997	AU 6835196 A CA 2227301 A EP 0852586 A	27-03-1997 13-03-1997 15-07-1998
EP 0318216	A	31-05-1989	AU 2796789 A CN 1035679 A CN 1073719 A DD 287104 A DD 298524 A DD 298525 A DD 298526 A DD 298527 A DE 3886363 D DE 3886363 T DK 353789 A ES 2012739 T FI 893447 A FI 981380 A GB 2212511 A,B GR 90300069 T HK 38293 A HR 940493 A HU 9500427 A IE 62868 B IN 171237 A IN 171238 A IN 171239 A IN 171240 A IN 169067 A JP 2810032 B JP 10108674 A JP 10290696 A JP 10290697 A JP 2662358 B JP 6225797 A JP 2810022 B JP 9184844 A JP 9173079 A JP 2809956 B JP 6128290 A JP 2662350 B JP 6128291 A JP 2532805 B JP 6128292 A JP 2662351 B JP 6128293 A JP 2500880 T LV 10726 A LV 10726 B NO 955101 A NO 955102 A PT 89041 A,B SI 8812138 A US 5714596 A	14-06-1989 20-09-1989 30-06-1993 14-02-1991 27-02-1992 27-02-1992 27-02-1992 27-02-1992 27-01-1994 19-05-1994 18-07-1989 01-12-1994 17-07-1989 15-06-1998 26-07-1989 31-07-1991 30-04-1993 31-08-1997 28-12-1995 08-03-1995 22-08-1992 22-08-1992 22-08-1992 22-08-1992 31-08-1991 15-10-1998 28-04-1998 04-11-1998 04-11-1998 08-10-1997 16-08-1994 15-10-1998 15-07-1997 08-07-1997 15-10-1998 10-05-1994 08-10-1997 10-05-1994 11-09-1996 10-05-1994 08-10-1997 10-05-1994 01-12-1988 31-08-1997 03-02-1998
EP 0388232	A	19-09-1990	AU 4750493 A AU 666767 B AU 666576 B	20-01-1994 22-02-1996 15-02-1996

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 98/01628

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0388232 A		AU 640920 B	09-09-1993
		AU 5278390 A	22-10-1990
		CA 2012482 A	17-09-1990
		DD 297446 A	09-01-1992
		FI 981381 A	15-06-1998
		GR 91300016 T	15-11-1991
		HU 9500746 A	29-01-1996
		JP 2693908 B	24-12-1997
		JP 7101986 A	18-04-1995
		JP 2809987 B	15-10-1998
		JP 7145194 A	06-06-1995
		JP 10309197 A	24-11-1998
		JP 10295389 A	10-11-1998
		JP 9215497 A	19-08-1997
		JP 2656995 B	24-09-1997
		JP 4504715 T	20-08-1992
		NO 303941 B	28-09-1998
		NO 960741 A	23-02-1996
		NO 960742 A	23-02-1996
		PT 93480 A, B	07-11-1990
		US 5714596 A	03-02-1998
		WO 9011089 A	04-10-1990
		US 5679342 A	21-10-1997
		US 5350671 A	27-09-1994
		US 5698390 A	16-12-1997
		US 5712088 A	27-01-1998
		US 5863719 A	26-01-1999
		US 5683864 A	04-11-1997
WO 9605513 A	22-02-1996	AU 3189495 A	07-03-1996
		CA 2174212 A	22-02-1996
		EP 0723665 A	31-07-1996
		JP 9504377 T	28-04-1997

PATENT COOPERATION TREATY

PCT

REC'D 05 JAN 2000

WIPO

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference <b>P18098WO</b>	<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. <b>PCT/IB98/01628</b>	International filing date (day/month/year) <b>06/10/1998</b>	Priority date (day/month/year) <b>06/10/1997</b>
International Patent Classification (IPC) or national classification and IPC <b>C12N15/12</b>		
Applicant <b>CHIRON S.P.A. et al.</b>		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 11 sheets, including this cover sheet.

☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☒ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand  <b>06/05/1999</b>	Date of completion of this report  <b>21.12.99</b>
Name and mailing address of the international preliminary examining authority:   <b>European Patent Office</b> <b>D-80298 Munich</b> <b>Tel. +49 89 2399 - 0 Tx: 523656 epmu d</b> <b>Fax: +49 89 2399 - 4465</b>	Authorized officer  <b>Julia, P</b>  Telephone No. <b>+49 89 2399 8410</b>  

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/IB98/01628

## I. Basis of the report

1. This report has been drawn on the basis of (*substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments.*):

### Description, pages:

1-28 as originally filed

### Claims, No.:

1-26 as originally filed

### Drawings, sheets:

1/17-17/17 as originally filed

2. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:
- ☐ the drawings, sheets:

3. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

4. Additional observations, if necessary:

**see separate sheet**

## II. Priority

1. ☐ This report has been established as if no priority had been claimed due to the failure to furnish within the prescribed time limit the requested:
- ☐ copy of the earlier application whose priority has been claimed.
  - ☐ translation of the earlier application whose priority has been claimed.
2. ☐ This report has been established as if no priority had been claimed due to the fact that the priority claim has

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/IB98/01628

been found invalid.

Thus for the purposes of this report, the international filing date indicated above is considered to be the relevant date.

3. Additional observations, if necessary:

**see separate sheet**

## V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

### 1. Statement

Novelty (N)	Yes:	Claims	19-21
	No:	Claims	1-18, 22-26
Inventive step (IS)	Yes:	Claims	
	No:	Claims	1-26
Industrial applicability (IA)	Yes:	Claims	1-26
	No:	Claims	

### 2. Citations and explanations

**see separate sheet**

## VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:

**see separate sheet**

## VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:

**see separate sheet**

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/IB98/01628

**1. Additional remarks to item I :**

A "Sequence Listing" has been filed by the Applicant on 28.02.99. This "Sequence Listing" comprises SEQ ID No.: 1 to SEQ ID No.: 15 (pages 1-6).

**2. Additional remarks to item II :**

The priority documents pertaining to the present application were not available at the time of establishing this international preliminary examination report (IPER). Hence, the current assessment is based on the assumption that all claims enjoy priority rights from the filing date of the priority document (06.10.97).

**3. Additional remarks to item V :**

The present application discloses that the DNA encoding the hepatitis C virus (HCV) receptor actually encodes the cellular protein known as CD81 (figure 1). This CD81 protein had been previously identified by monoclonal antibodies as the target of an antiproliferative antibody (TAPA-1) which inhibited in vitro cellular proliferation. The "EC2" region of CD81 (extracellular loop 2, second extracellular region of CD81) (residues 113-201) is the region involved in binding to the E2 protein of HCV. The application discloses the CD81 protein of chimpanzee, green monkey, hamster, rat and mouse too (figure 1). The application explicitly claims compounds that bind specifically to a CD81 protein (for use in the therapy or diagnosis of HCV), methods for screening such compounds, methods of treating, pharmaceutical compositions, uses in the manufacture of medicaments, assays for detecting HCV antibodies or HCV in a serum sample, diagnostic kits, transgenic non-human mammals, etc...

The following documents have been cited in the International Search Report (ISR) as being relevant for assessing the novelty and inventiveness of the claimed subject matter:

i) R. Oren et al. , Mol. Cell. Biol. 1990, Vol. 10 (8), pages 4007-4015 (**D1**) discloses the identification, cloning and characterization of the human 26-kd cell surface protein TAPA-1 (CD81) using a murine (cell) antiproliferative monoclonal antibody. The cDNA sequence and the corresponding deduced amino acid sequence are shown in figure 3. Reference is also made to an hybridization probe (coding and 3' untranslated region) used in genomic Southern blots for probing enzymatically digested genomic DNA. The deduced TAPA-1 amino acid sequence is said to be highly homologous to the leukocyte membrane protein CD37 and to the melanoma-associated antigen ME491. There is, however, no reference



in D1 to the possible use of the TAPA-1 antigen (CD81 protein) in the therapy or diagnosis of HCV.

ii) a similar disclosure is found in document M.L. Andria et al., J. Immunol. 1991, Vol. 147 (3), pages 1030-1036 (**D2**) which refers to the genomic organization and chromosomal localization of the TAPA-1 gene. D2 also refers to the homology of TAPA-1 to additional proteins, namely a cell surface marker on both haematopoietic and non-haematopoietic tissues (CD9), a pan-leukocyte antigen (CD53), a colon cancer antigen (CO-029) and a schistosome surface antigen (Sm23). Figure 3 outlines the relevance of the second extracellular loop of TAPA-1. There is, however, no reference in D2 to the possible use of the TAPA-1 antigen (CD81 protein) in the therapy or diagnosis of HCV.

In view of D1 and D2, the IPEA considers that the particular intended use disclosed in the application (i.e. therapy or diagnosis of HCV) must be seen as a distinctive characteristic of the claimed product which can not be disregarded for assessing the novelty and inventiveness of said claimed product (PCT Gazette-Section IV, Special Issue 29.10.98; PCT International Preliminary Examination Guidelines, Chapter III-4.8 and in particular Chapter IV-7.6) (first medical application). However, the IPEA considers that :

a) claim 4 is directed to a protein **comprising** the second extracellular region of CD81, i.e. the claim is not limited to this second extracellular region but it embraces any protein comprising said region and it is further not restricted to any intended use (therapy or diagnosis of HCV). The IPEA considers that the known TAPA-1 protein (CD81) certainly comprises this second extracellular loop and thus, D1 and D2 clearly anticipate the subject matter of claim 4 (Articles 33 (2) and (3) PCT).

b) claims 22-24 are directed to general nucleic acid molecules hybridizing with a nucleic acid molecule which encodes a CD81 protein. In view of the nucleotide sequences and the specific probes disclosed in D1 and D2, the IPEA considers that this subject matter is clearly anticipated by those documents (Articles 33 (2) and (3) PCT).

c) claims 8, 9 and 11 are directed to a general pharmaceutical composition comprising a CD81 protein or a protein comprising the second extracellular region of the CD81 protein and to a method for preparing said pharmaceutical compositions (there are no references to HCV). Claims 16-17 are directed to a general diagnostic kit comprising a CD81 protein

but without any reference to HCV. In view of the high homology to well-known proteins and/or antigens which had already been used in general pharmaceutical compositions, diagnostic kits, etc... the IPEA considers that the skilled person would certainly have thought in using the TAPA-1 proteins in similar compositions. Thus, these claims do not fulfil the requirements of Article 33 (3) PCT.

d) claims 19 and 20 are directed to a transgenic non-human mammal carrying a transgene encoding CD81 protein and a process for producing such a transgenic animal. The IPEA considers that once the nucleotide sequence of TAPA-1 was known and available to the skilled person, the production of a transgenic animal (mouse) would not have required any inventive contribution from the skilled person (Article 33 (3) PCT).

iii) WO-A-97/09349 (D3) discloses a human transmembrane cellular protein (putative cellular receptor) of 24-kD which is capable of specifically binding the E2 envelope protein of HCV. D3 further refers to pharmaceutical compositions comprising said protein, uses thereof for HCV diagnosis and therapy, methods for preparing such protein and for treating HCV, assay for detecting HCV antibodies, diagnostic kits, transgenic animals, etc... However, D3 actually discloses only the isolation of the 24-kD protein and its partial (functional) characterization but there is no amino acid sequence let alone a nucleotide sequence in D3.

In view of the (functional) characterization disclosed in D3, the IPEA considers that the protein isolated in D3 is the CD81 protein and that the actual contribution of the present application is the determination of its amino acid sequence and the nucleotide sequence of the corresponding gene and/or cDNA. However, the IPEA considers that once a protein is known and available to the skilled person, the determination of an inherent feature such as the amino acid sequence does not change any property of the known product per se and thus, it can not render novel this product. In addition, a document disclosing a (chemical) compound and its isolation and/or manufacture makes normally available this compound to the public in all desired degrees of purity (unless exceptional situations wherein all conventional purification processes have failed, etc...). The attention of the Applicant is also drawn to the fact that irrespective whether the protein disclosed in D3 is actually the claimed CD81 protein or not, the protein disclosed in D3 is certainly a "functional equivalent" of the CD81 protein (as it binds to the E2 of HCV). Thus, claims directed to "a CD81 protein or a functional equivalent thereof", namely pharmaceutical

compositions, methods and uses, etc... as well as claims directed to a compound that binds specifically to a CD81 protein, namely the subject matter of claims 1-18 and 25-26 do not fulfil the requirements of Articles 33 (2) and (3) PCT.

Furthermore, once a protein has been identified, the determination of a nucleic acid coding said protein does not require any specific inventive skill, i.e. at the priority date a skilled person could expect to perform the cloning and expression of a gene in a fairly straightforward manner, and if the cloning, although requiring much work, does not pose such problems as to prove that the expectation of success was ill-founded, then inventive step can not be acknowledged. Thus, the IPEA considers that the subject matter of claims 19-24 does not fulfil the requirements of Article 33 (3) PCT over the teachings of D3.

iv) documents EP-A-0 318 216 (**D4**) and EP-A-0 388 232 (**D5**) are concerned with the characterization of the HCV, wherein the nucleotide sequence and amino acid sequence of the envelope region of the HCV (comprising the E2 region) are explicitly disclosed. Similarly the document D. Rosa et al., Proc. Natl. Acad. Sci. USA 1996, Vol. 93, pages 1759-1763 (**D6**) outlines the importance of the E2 region of HCV for target cellular binding. In this respect, WO-A-96/05513 (**D7**) emphasizes too the importance of the E2 region of HCV and its use for the detecting and identifying the cellular receptor for HCV. Thus, these documents are considered to anticipate at least the subject matter of claim 6 (Articles 33 (2) and (3) PCT).

The attention of the Applicant is also drawn to the subject matter of claim 7 which is directed to a method of treatment of the human and/or animal body and thus, it may be excluded from examination by Article 34(4)(a)(i) PCT in combination with Rule 67(iv) PCT too. Furthermore, for such a subject matter no unified criteria exist in PCT for the assessment whether it is industrially applicable or not. The patentability can also be dependent upon the formulation of the claims. The EPO, for example, does not recognize as industrially applicable the subject matter of claims to the use of a compound in medical treatment, but will allow, however, claims to a known compound for first use in medical treatment (such as present claims 1-3, 5-6, etc...) and the use of such a compound for the manufacture of a medicament for a new medical treatment (claims 12-13).

**4. Additional remarks to item VII :**

According to the ISA the Accession number of the CD81 protein in EMBL/GENBANK database is wrong. Accession number M33690 corresponds to rat alpha 2U globulin promoter region, whereas M33680 actually corresponds to human CD81.

**5. Additional remarks to item VIII :**

The following objections are raised under **Article 6 PCT** concerning the clarity of the claims :

i) according to Article 6 PCT in combination with Rule 6.3 PCT the claims shall define the matter for which protection is sought in terms of technical features. The IPEA considers that a peptide, polypeptide, protein, oligonucleotide, gene, etc.. being chemical products must be clearly and unambiguously characterized by their amino acid and/or nucleic acid sequences, i.e. by reference to their specific SEQ ID No or Figure(s) comprising said sequences. The characterization of a product only by the desired function or by an arbitrary abbreviation without any real technical meaning (CD81) does not fulfill the requirements of said Article 6 PCT in combination with Rule 6.3 PCT. Thus, this objection applies for the subject matter of claims 1, 6-8, 10-12 and 14-26.

ii) the IPEA considers that the general reference to a "functional equivalent thereof" is ambiguous as far as said function is not clearly defined in the wording of the claim. A protein can have many "functions" depending on its intended use, such as for instance as a substrate and/or inhibitor of particular enzymes (protease, methylases, etc...), an immunogen, foodstuff additive, etc... Even if the claim refers to the particular "use in the therapy or diagnosis of HCV", the IPEA considers that such a use per se does not clearly and unambiguously characterize the meaning of the wording "functional equivalent", in particular in view of the definition found in the description, namely "any fragment or assembly of fragments of the complete protein that binds to HCV" (page 3 lines 7-9) (under suitable conditions (pH, salt, temperature, etc...) any peptide can bind to another polypeptide and wherein said peptide certainly comprises an arbitrarily short fragment (one, two, three residues) of any other peptide too) (the description refers to an homology of only 25% too !!). Thus, the scope of claims 1-2, 4, 7-8, 11-12, 14-16, 18-19, 21 and 25-26 is ambiguous.

iii) in this respect the application discloses the presence and the sequence of CD81

proteins in hamster, rat and mouse (figure 1). These CD81 sequences are closely related, and they have a high homology (greater than 80%, with fragments being 100% homologous) to the CD81 sequences of human and chimpanzee. In particular, the CD81 protein from hamster has only 10 exchanged in the EC2 region (89 residues), the rat CD81 14 exchanged residues and the mouse CD81 only 17 residues (the Applicant is reminded, however, that most of the claims are directed to "a CD81 protein" and not to the specific "human and/or chimpanzee CD81 E2, known to be responsible for binding the HCV). As the Applicant rightly emphasizes, the "HCV receptor, whilst ubiquitous in humans and found in chimpanzees, is absent in other mammals" (page 12 lines 21-22). Thus, the IPEA considers that (a) whereas the CD81 proteins of these animals and fragments thereof are embraced by the general wording "a CD81 protein" and they are certainly "functional equivalents" of a CD81 protein, these "CD81 proteins" are not suitable, however, for the intended use (therapy or diagnostic of HCV). Furthermore, (b) transgenic animals comprising such "CD81 proteins" do not seem to be useful as "animal models of HCV infection" (see also example 7 on page 24, wherein it is also explicitly acknowledged that "human, but not the mouse EC2 loop of CD81, bound to E2 ... and inhibited binding of E2 to human cells").

iv) whereas there is no clarity problems for a claim directed to "a method for screening chemical compounds for ability to bind to the region of HCV responsible for binding to a host cell" wherein said method uses the CD81 protein (claim 18), clarity problems arise for claims directed to such products "per se". Claim 6 is only characterized by the result to be achieved, namely "capable of binding specifically to a CD81 protein". The claimed product is only characterized by the desired functional property (desiratum) but not by any actual technical feature (parameters, structure, sequence, etc...). As far as said products are not clearly and unambiguously characterized, the IPEA considers that such a claim could embrace well-known products which under certain suitable conditions (pH, salt, temperature, etc...) could "specifically bind" the CD81 protein (the claim is not even limited to a protein, peptide and/or polypeptide but embraces any possible "compound"). The same objection applies too to the subject matter of claims 7-8 and 10-12.

v) in this respect, the description refers to the "ideal" compound that binds to CD81 as the one which only interacts with CD81 in the presence of HCV, so that the usual function of CD81 is not compromised on healthy cells. This requirement is not present in the wording of any claim related to such products, thus it is not clear whether a "compound" as claimed

could be actually used "in the therapy or diagnosis of HCV" (claims 6-7, 10 and 12). In addition, it is not clear whether such "ideal" compounds could be easily screened by the method of claim 18, which only requires the presence of the chemical compound to be screened and the CD81 protein but in the absence of HCV.

vi) contrary to the subject matter of claim 15 which is directed to "an assay for HCV ", the subject matter of claim 14 is directed to an assay for general "HCV antibodies". However, the reference to "HCV protein" in claim 14 is ambiguous. There are many "HCV proteins" (produced by proteolysis of the HCV polyprotein) and it is not clear from said general wording whether all of them could be successfully used in the claimed assay or alternatively all of them should be used in the claimed assay.

vii) the subject matter of claims 19 and 20 refers to general "non-human transgenic mammals". The IPEA considers that such a broad subject matter is not fully (technically) supported by the present description. Methods for producing mammalian animals are not general for all mammals and they depend on the specific mammal used. Several steps such as a suitable status (stage) of the egg for optimal introduction of the DNA construct(s), time of insertion of the egg into the oviduct or uterus, etc.. are critical for the success of the method. Furthermore, several mammalian species do not lend themselves applicable to the present technology used for transgenic mice (or sheep, rabbits, goats and cows), such as duckbill platypus, spiny anteaters (echidna, egg-laying mammals, etc..). The IPEA considers that the description is not enabling for whole range claimed, i.e. does not allow the skilled person to produce the claimed product in or for the whole range claimed (all mammalian species). The specification only teaches the production of a single mammalian species, namely mouse. It would be highly unpredictable and would require undue experimentation to prepare a homologous population of all mammalian species. The preparation of one species is not sufficient evidence to support that it can be prepared from all mammalian species.

viii) the wording of claim 22 is completely ambiguous. There is neither a functional requirement nor a minimum length required for the claimed nucleic acid molecule. Furthermore, the claim does not provide any clear definition or characterization of the cited "standard conditions", which thus are completely ambiguous. In fact, the scope of the claim embraces any (short) arbitrary nucleic acid (by selecting suitable "standard conditions") without bearing any technical relationship to the disclosed CD81 protein). The first part of

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/IB98/01628

this objection (function and length) also applies for the subject matter of claim 23 (and dependent claim 24).

From the  
INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

PCT

NOTIFICATION OF TRANSMITTAL OF  
THE INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT  
(PCT Rule 71.1)

To:

HALLYBONE, Huw George  
CARPMAELS & RANSFORD  
43 Bloomsbury Square  
London WC1A 2RA  
GRANDE BRETAGNE

Date of mailing  
(day/month/year)

21.12.99

Applicant's or agent's file reference  
P18098WO

IMPORTANT NOTIFICATION

International application No.  
PCT/IB98/01628

International filing date (day/month/year)  
06/10/1998

Priority date (day/month/year)  
06/10/1997

Applicant  
CHIRON S.P.A. et al.

1. The applicant is hereby notified that this International Preliminary Examining Authority transmits herewith the international preliminary examination report and its annexes, if any, established on the international application.
2. A copy of the report and its annexes, if any, is being transmitted to the International Bureau for communication to all the elected Offices.
3. Where required by any of the elected Offices, the International Bureau will prepare an English translation of the report (but not of any annexes) and will transmit such translation to those Offices.

4. REMINDER

The applicant must enter the national phase before each elected Office by performing certain acts (filing translations and paying national fees) within 30 months from the priority date (or later in some Offices) (Article 39(1)) (see also the reminder sent by the International Bureau with Form PCT/IB/301).

Where a translation of the international application must be furnished to an elected Office, that translation must contain a translation of any annexes to the international preliminary examination report. It is the applicant's responsibility to prepare and furnish such translation directly to each elected Office concerned.

For further details on the applicable time limits and requirements of the elected Offices, see Volume II of the PCT Applicant's Guide.

Name and mailing address of the IPEA/



European Patent Office  
D-80298 Munich  
Tel. +49 89 2399 - 0 Tx: 523656 epmu d  
Fax: +49 89 2399 - 4465

Authorized officer

Schou, S

Tel. +49 89 2399-8718





# PATENT COOPERATION TREATY

## PCT

### INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)



Applicant's or agent's file reference <b>P18098WO</b>	<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. <b>PCT/IB98/01628</b>	International filing date (day/month/year) <b>06/10/1998</b>	Priority date (day/month/year) <b>06/10/1997</b>
International Patent Classification (IPC) or national classification and IPC <b>C12N15/12</b>		
Applicant <b>CHIRON S.P.A. et al.</b>		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2. This REPORT consists of a total of 11 sheets, including this cover sheet.  
  
☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☒ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand  <b>06/05/1999</b>	Date of completion of this report  <b>21.12.99</b>
Name and mailing address of the international preliminary examining authority:   <b>European Patent Office</b> <b>D-80298 Munich</b> <b>Tel. +49 89 2399 - 0 Tx: 523656 epmu d</b> <b>Fax: +49 89 2399 - 4465</b>	Authorized officer  <b>Julia, P</b>   <b>Telephone No. +49 89 2399 8410</b>

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT**

International application No. PCT/IB98/01628

**I. Basis of the report**

1. This report has been drawn on the basis of (*substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments.*):

**Description, pages:**

1-28 as originally filed

**Claims, No.:**

1-26 as originally filed

**Drawings, sheets:**

1/17-17/17 as originally filed

2. The amendments have resulted in the cancellation of:

- ☐ the description, pages:  
☐ the claims, Nos.:  
☐ the drawings, sheets:

3. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

4. Additional observations, if necessary:

**see separate sheet**

**II. Priority**

1. ☐ This report has been established as if no priority had been claimed due to the failure to furnish within the prescribed time limit the requested:
- ☐ copy of the earlier application whose priority has been claimed.
- ☐ translation of the earlier application whose priority has been claimed.
2. ☐ This report has been established as if no priority had been claimed due to the fact that the priority claim has

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT**

International application No. PCT/IB98/01628

been found invalid.

Thus for the purposes of this report, the international filing date indicated above is considered to be the relevant date.

3. Additional observations, if necessary:

**see separate sheet**

**V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**

**1. Statement**

Novelty (N)	Yes:	Claims	19-21
	No:	Claims	1-18, 22-26
Inventive step (IS)	Yes:	Claims	
	No:	Claims	1-26
Industrial applicability (IA)	Yes:	Claims	1-26
	No:	Claims	

**2. Citations and explanations**

**see separate sheet**

**VII. Certain defects in the international application**

The following defects in the form or contents of the international application have been noted:

**see separate sheet**

**VIII. Certain observations on the international application**

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:

**see separate sheet**

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/IB98/01628

**1. Additional remarks to item I :**

A "Sequence Listing" has been filed by the Applicant on 28.02.99. This "Sequence Listing" comprises SEQ ID No.: 1 to SEQ ID No.: 15 (pages 1-6).

**2. Additional remarks to item II :**

The priority documents pertaining to the present application were not available at the time of establishing this international preliminary examination report (IPER). Hence, the current assessment is based on the assumption that all claims enjoy priority rights from the filing date of the priority document (06.10.97).

**3. Additional remarks to item V :**

The present application discloses that the DNA encoding the hepatitis C virus (HCV) receptor actually encodes the cellular protein known as CD81 (figure 1). This CD81 protein had been previously identified by monoclonal antibodies as the target of an antiproliferative antibody (TAPA-1) which inhibited in vitro cellular proliferation. The "EC2" region of CD81 (extracellular loop 2, second extracellular region of CD81) (residues 113-201) is the region involved in binding to the E2 protein of HCV. The application discloses the CD81 protein of chimpanzee, green monkey, hamster, rat and mouse too (figure 1). The application explicitly claims compounds that bind specifically to a CD81 protein (for use in the therapy or diagnosis of HCV), methods for screening such compounds, methods of treating, pharmaceutical compositions, uses in the manufacture of medicaments, assays for detecting HCV antibodies or HCV in a serum sample, diagnostic kits, transgenic non-human mammals, etc...

The following documents have been cited in the International Search Report (ISR) as being relevant for assessing the novelty and inventiveness of the claimed subject matter:

i) R. Oren et al. , Mol. Cell. Biol. 1990, Vol. 10 (8), pages 4007-4015 (D1) discloses the identification, cloning and characterization of the human 26-kd cell surface protein TAPA-1 (CD81) using a murine (cell) antiproliferative monoclonal antibody. The cDNA sequence and the corresponding deduced amino acid sequence are shown in figure 3. Reference is also made to an hybridization probe (coding and 3' untranslated region) used in genomic Southern blots for probing enzymatically digested genomic DNA. The deduced TAPA-1 amino acid sequence is said to be highly homologous to the leukocyte membrane protein CD37 and to the melanoma-associated antigen ME491. There is, however, no reference

in D1 to the possible use of the TAPA-1 antigen (CD81 protein) in the therapy or diagnosis of HCV.

ii) a similar disclosure is found in document M.L. Andria et al., J. Immunol. 1991, Vol. 147 (3), pages 1030-1036 (D2) which refers to the genomic organization and chromosomal localization of the TAPA-1 gene. D2 also refers to the homology of TAPA-1 to additional proteins, namely a cell surface marker on both haematopoietic and non-haematopoietic tissues (CD9), a pan-leukocyte antigen (CD53), a colon cancer antigen (CO-029) and a schistosome surface antigen (Sm23). Figure 3 outlines the relevance of the second extracellular loop of TAPA-1. There is, however, no reference in D2 to the possible use of the TAPA-1 antigen (CD81 protein) in the therapy or diagnosis of HCV.

In view of D1 and D2, the IPEA considers that the particular intended use disclosed in the application (i.e. therapy or diagnosis of HCV) must be seen as a distinctive characteristic of the claimed product which can not be disregarded for assessing the novelty and inventiveness of said claimed product (PCT Gazette-Section IV, Special Issue 29.10.98; PCT International Preliminary Examination Guidelines, Chapter III-4.8 and in particular Chapter IV-7.6) (first medical application). However, the IPEA considers that :

a) claim 4 is directed to a protein **comprising** the second extracellular region of CD81, i.e. the claim is not limited to this second extracellular region but it embraces any protein comprising said region and it is further not restricted to any intended use (therapy or diagnosis of HCV). The IPEA considers that the known TAPA-1 protein (CD81) certainly comprises this second extracellular loop and thus, D1 and D2 clearly anticipate the subject matter of claim 4 (Articles 33 (2) and (3) PCT).

b) claims 22-24 are directed to general nucleic acid molecules hybridizing with a nucleic acid molecule which encodes a CD81 protein. In view of the nucleotide sequences and the specific probes disclosed in D1 and D2, the IPEA considers that this subject matter is clearly anticipated by those documents (Articles 33 (2) and (3) PCT).

c) claims 8, 9 and 11 are directed to a general pharmaceutical composition comprising a CD81 protein or a protein comprising the second extracellular region of the CD81 protein and to a method for preparing said pharmaceutical compositions (there are no references to HCV). Claims 16-17 are directed to a general diagnostic kit comprising a CD81 protein

but without any reference to HCV. In view of the high homology to well-known proteins and/or antigens which had already been used in general pharmaceutical compositions, diagnostic kits, etc... the IPEA considers that the skilled person would certainly have thought in using the TAPA-1 proteins in similar compositions. Thus, these claims do not fulfil the requirements of Article 33 (3) PCT.

d) claims 19 and 20 are directed to a transgenic non-human mammal carrying a transgene encoding CD81 protein and a process for producing such a transgenic animal. The IPEA considers that once the nucleotide sequence of TAPA-1 was known and available to the skilled person, the production of a transgenic animal (mouse) would not have required any inventive contribution from the skilled person (Article 33 (3) PCT).

iii) WO-A-97/09349 (D3) discloses a human transmembrane cellular protein (putative cellular receptor) of 24-kD which is capable of specifically binding the E2 envelope protein of HCV. D3 further refers to pharmaceutical compositions comprising said protein, uses thereof for HCV diagnosis and therapy, methods for preparing such protein and for treating HCV, assay for detecting HCV antibodies, diagnostic kits, transgenic animals, etc... However, D3 actually discloses only the isolation of the 24-kD protein and its partial (functional) characterization but there is no amino acid sequence let alone a nucleotide sequence in D3.

In view of the (functional) characterization disclosed in D3, the IPEA considers that the protein isolated in D3 is the CD81 protein and that the actual contribution of the present application is the determination of its amino acid sequence and the nucleotide sequence of the corresponding gene and/or cDNA. However, the IPEA considers that once a protein is known and available to the skilled person, the determination of an inherent feature such as the amino acid sequence does not change any property of the known product per se and thus, it can not render novel this product. In addition, a document disclosing a (chemical) compound and its isolation and/or manufacture makes normally available this compound to the public in all desired degrees of purity (unless exceptional situations wherein all conventional purification processes have failed, etc...). The attention of the Applicant is also drawn to the fact that irrespective whether the protein disclosed in D3 is actually the claimed CD81 protein or not, the protein disclosed in D3 is certainly a "functional equivalent" of the CD81 protein (as it binds to the E2 of HCV). Thus, claims directed to "a CD81 protein or a functional equivalent thereof", namely pharmaceutical

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/IB98/01628

compositions, methods and uses, etc... as well as claims directed to a compound that binds specifically to a CD81 protein, namely the subject matter of claims 1-18 and 25-26 do not fulfil the requirements of Articles 33 (2) and (3) PCT.

Furthermore, once a protein has been identified, the determination of a nucleic acid coding said protein does not require any specific inventive skill, i.e. at the priority date a skilled person could expect to perform the cloning and expression of a gene in a fairly straightforward manner, and if the cloning, although requiring much work, does not pose such problems as to prove that the expectation of success was ill-founded, then inventive step can not be acknowledged. Thus, the IPEA considers that the subject matter of claims 19-24 does not fulfil the requirements of Article 33 (3) PCT over the teachings of D3.

iv) documents EP-A-0 318 216 (D4) and EP-A-0 388 232 (D5) are concerned with the characterization of the HCV, wherein the nucleotide sequence and amino acid sequence of the envelope region of the HCV (comprising the E2 region) are explicitly disclosed. Similarly the document D. Rosa et al., Proc. Natl. Acad. Sci. USA 1996, Vol. 93, pages 1759-1763 (D6) outlines the importance of the E2 region of HCV for target cellular binding. In this respect, WO-A-96/05513 (D7) emphasizes too the importance of the E2 region of HCV and its use for the detecting and identifying the cellular receptor for HCV. Thus, these documents are considered to anticipate at least the subject matter of claim 6 (Articles 33 (2) and (3) PCT).

The attention of the Applicant is also drawn to the subject matter of claim 7 which is directed to a method of treatment of the human and/or animal body and thus, it may be excluded from examination by Article 34(4)(a)(i) PCT in combination with Rule 67(iv) PCT too. Furthermore, for such a subject matter no unified criteria exist in PCT for the assessment whether it is industrially applicable or not. The patentability can also be dependent upon the formulation of the claims. The EPO, for example, does not recognize as industrially applicable the subject matter of claims to the use of a compound in medical treatment, but will allow, however, claims to a known compound for first use in medical treatment (such as present claims 1-3, 5-6, etc...) and the use of such a compound for the manufacture of a medicament for a new medical treatment (claims 12-13).

**4. Additional remarks to item VII :**

According to the ISA the Accession number of the CD81 protein in EMBL/GENBANK database is wrong. Accession number M33690 corresponds to rat alpha 2U globulin promoter region, whereas M33680 actually corresponds to human CD81.

**5. Additional remarks to item VIII :**

The following objections are raised under **Article 6 PCT** concerning the clarity of the claims :

i) according to Article 6 PCT in combination with Rule 6.3 PCT the claims shall define the matter for which protection is sought in terms of technical features. The IPEA considers that a peptide, polypeptide, protein, oligonucleotide, gene, etc.. being chemical products must be clearly and unambiguously characterized by their amino acid and/or nucleic acid sequences, i.e. by reference to their specific SEQ ID No or Figure(s) comprising said sequences. The characterization of a product only by the desired function or by an arbitrary abbreviation without any real technical meaning (CD81) does not fulfill the requirements of said Article 6 PCT in combination with Rule 6.3 PCT. Thus, this objection applies for the subject matter of claims 1, 6-8, 10-12 and 14-26.

ii) the IPEA considers that the general reference to a "functional equivalent thereof" is ambiguous as far as said function is not clearly defined in the wording of the claim. A protein can have many "functions" depending on its intended use, such as for instance as a substrate and/or inhibitor of particular enzymes (protease, methylases, etc...), an immunogen, foodstuff additive, etc... Even if the claim refers to the particular "use in the therapy or diagnosis of HCV", the IPEA considers that such a use per se does not clearly and unambiguously characterize the meaning of the wording "functional equivalent", in particular in view of the definition found in the description, namely "any fragment or assembly of fragments of the complete protein that binds to HCV" (page 3 lines 7-9) (under suitable conditions (pH, salt, temperature, etc...) any peptide can bind to another polypeptide and wherein said peptide certainly comprises an arbitrarily short fragment (one, two, three residues) of any other peptide too) (the description refers to an homology of only 25% too !!). Thus, the scope of claims 1-2, 4, 7-8, 11-12, 14-16, 18-19, 21 and 25-26 is ambiguous.

iii) in this respect the application discloses the presence and the sequence of CD81



**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/IB98/01628

proteins in hamster, rat and mouse (figure 1). These CD81 sequences are closely related, and they have a high homology (greater than 80%, with fragments being 100% homologous) to the CD81 sequences of human and chimpanzee. In particular, the CD81 protein from hamster has only 10 exchanged in the EC2 region (89 residues), the rat CD81 14 exchanged residues and the mouse CD81 only 17 residues (the Applicant is reminded, however, that most of the claims are directed to "a CD81 protein" and not to the specific "human and/or chimpanzee CD81 E2, known to be responsible for binding the HCV). As the Applicant rightly emphasizes, the "HCV receptor, whilst ubiquitous in humans and found in chimpanzees, is absent in other mammals" (page 12 lines 21-22). Thus, the IPEA considers that (a) whereas the CD81 proteins of these animals and fragments thereof are embraced by the general wording "a CD81 protein" and they are certainly "functional equivalents" of a CD81 protein, these "CD81 proteins" are not suitable, however, for the intended use (therapy or diagnostic of HCV). Furthermore, (b) transgenic animals comprising such "CD81 proteins" do not seem to be useful as "animal models of HCV infection" (see also example 7 on page 24, wherein it is also explicitly acknowledged that "human, but not the mouse EC2 loop of CD81, bound to E2 ... and inhibited binding of E2 to human cells").

iv) whereas there is no clarity problems for a claim directed to "a method for screening chemical compounds for ability to bind to the region of HCV responsible for binding to a host cell" wherein said method uses the CD81 protein (claim 18), clarity problems arise for claims directed to such products "per se". Claim 6 is only characterized by the result to be achieved, namely "capable of binding specifically to a CD81 protein". The claimed product is only characterized by the desired functional property (desiratum) but not by any actual technical feature (parameters, structure, sequence, etc...). As far as said products are not clearly and unambiguously characterized, the IPEA considers that such a claim could embrace well-known products which under certain suitable conditions (pH, salt, temperature, etc...) could "specifically bind" the CD81 protein (the claim is not even limited to a protein, peptide and/or polypeptide but embraces any possible "compound"). The same objection applies too to the subject matter of claims 7-8 and 10-12.

v) in this respect, the description refers to the "ideal" compound that binds to CD81 as the one which only interacts with CD81 in the presence of HCV, so that the usual function of CD81 is not compromised on healthy cells. This requirement is not present in the wording of any claim related to such products, thus it is not clear whether a "compound" as claimed

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/IB98/01628

could be actually used "in the therapy or diagnosis of HCV" (claims 6-7, 10 and 12). In addition, it is not clear whether such "ideal" compounds could be easily screened by the method of claim 18, which only requires the presence of the chemical compound to be screened and the CD81 protein but in the absence of HCV.

vi) contrary to the subject matter of claim 15 which is directed to "an assay for HCV", the subject matter of claim 14 is directed to an assay for general "HCV antibodies". However, the reference to "HCV protein" in claim 14 is ambiguous. There are many "HCV proteins" (produced by proteolysis of the HCV polyprotein) and it is not clear from said general wording whether all of them could be successfully used in the claimed assay or alternatively all of them should be used in the claimed assay.

vii) the subject matter of claims 19 and 20 refers to general "non-human transgenic mammals". The IPEA considers that such a broad subject matter is not fully (technically) supported by the present description. Methods for producing mammalian animals are not general for all mammals and they depend on the specific mammal used. Several steps such as a suitable status (stage) of the egg for optimal introduction of the DNA construct(s), time of insertion of the egg into the oviduct or uterus, etc.. are critical for the success of the method. Furthermore, several mammalian species do not lend themselves applicable to the present technology used for transgenic mice (or sheep, rabbits, goats and cows), such as duckbill platypus, spiny anteaters (echidna, egg-laying mammals, etc..). The IPEA considers that the description is not enabling for whole range claimed, i.e. does not allow the skilled person to produce the claimed product in or for the whole range claimed (all mammalian species). The specification only teaches the production of a single mammalian species, namely mouse. It would be highly unpredictable and would require undue experimentation to prepare a homologous population of all mammalian species. The preparation of one species is not sufficient evidence to support that it can be prepared from all mammalian species.

viii) the wording of claim 22 is completely ambiguous. There is neither a functional requirement nor a minimum length required for the claimed nucleic acid molecule. Furthermore, the claim does not provide any clear definition or characterization of the cited "standard conditions", which thus are completely ambiguous. In fact, the scope of the claim embraces any (short) arbitrary nucleic acid (by selecting suitable "standard conditions") without bearing any technical relationship to the disclosed CD81 protein). The first part of

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/IB98/01628

this objection (function and length) also applies for the subject matter of claim 23 (and dependent claim 24).

# PCT

## INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference <b>P18098W0</b>	<b>FOR FURTHER ACTION</b> see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. <b>PCT/IB 98/ 01628</b>	International filing date (day/month/year) <b>06/10/1998</b>	(Earliest) Priority Date (day/month/year) <b>06/10/1997</b>
Applicant <b>CHIRON S.P.A. et al.</b>		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

- ☒ Certain claims were found unsearchable (see Box I).
- ☐ Unity of invention is lacking (see Box II).
- ☒ The international application contains disclosure of a nucleotide and/or amino acid sequence listing and the international search was carried out on the basis of the sequence listing

☐ filed with the international application.

☒ furnished by the applicant separately from the international application,

☐ but not accompanied by a statement to the effect that it did not include matter going beyond the disclosure in the international application as filed.

☐ Transcribed by this Authority

- With regard to the title, ☐ the text is approved as submitted by the applicant  
☒ the text has been established by this Authority to read as follows:

**HEPATITIS C RECEPTOR PROTEIN CD81**

- With regard to the abstract, ☒ the text is approved as submitted by the applicant  
☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this International Search Report, submit comments to this Authority.

- The figure of the drawings to be published with the abstract is:

Figure No. 1A ☐ as suggested by the applicant.

☐ None of the figures.

☒ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB 98/01628

## Box I Observations where certain claims were found uns archable (Continuation f item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 7 and 20  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/IB 98/01628

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6    C12N15/12    C12N15/62    C07K14/705    A61K38/17    A01K67/027 G01N33/50    G01N33/576		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6    C12N    C07K    A61K    A01K    G01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	OREN R ET AL: "TAPA-1, THE TARGET OF AN ANTIPROLIFERATIVE ANTIBODY, DEFINES A NEW FAMILY OF TRANSMEMBRANE PROTEINS" MOLECULAR AND CELLULAR BIOLOGY, vol. 10, no. 8, August 1990, pages 4007-4015, XP000749300 see the whole document ---	1-5, 21-25
X	ANDRIA M L ET AL: "GENOMIC ORGANIZATION AND CHROMOSOMAL LOCALIZATION OF THE TAPA-1 GENE" JOURNAL OF IMMUNOLOGY, vol. 147, no. 3, 1 August 1991, pages 1030-1036, XP002064247 see the whole document --- <div style="text-align: center;">-/--</div>	1-5, 21-25
<div style="display: flex; justify-content: space-between;"> <span><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</span> <span><input checked="" type="checkbox"/> Patent family members are listed in annex.</span> </div>		
° Special categories of cited documents : <div style="display: flex;"> <div style="flex: 1;"> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="flex: 1;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&amp;" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search	Date of mailing of the international search report	
8 February 1999	18/02/1999	
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016	Authorized officer  Hornig, H	

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/IB 98/01628

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 09349 A (BIOCINE SPA ;ABRIGNANI SERGIO (IT)) 13 March 1997 cited in the application see the whole document ----	6
X	EP 0 318 216 A (CHIRON CORP) 31 May 1989 cited in the application see the whole document ----	6
X	EP 0 388 232 A (CHIRON CORP) 19 September 1990 cited in the application see the whole document ----	6
A	ROSA D ET AL: "A QUANTITATIVE TEST TO ESTIMATE NEUTRALIZING ANTIBODIES TO THE HEPATITIS C VIRUS: CYTOFLUORIMETRIC ASSESSMENT OF ENVELOPE GLYCOPROTEIN 2 BINDING TO TARGET CELLS" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 93, no. 3, March 1996, pages 1759-1763, XP000615446 cited in the application see the whole document ----	1-26
A	WO 96 05513 A (BIOCINE SPA ;ABRIGNANI SERGIO (IT)) 22 February 1996 cited in the application see the whole document ----	1-26
T	P. PILERI ET AL.: "Binding of hepatitis C virus to CD81" SCIENCE, vol. 282, 30 October 1998, pages 938-941, XP002092549 AAAS, WASHINGTON, DC, US see the whole document -----	1-26

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 98/01628

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9709349	A	13-03-1997	AU 6835196 A CA 2227301 A EP 0852586 A	27-03-1997 13-03-1997 15-07-1998
EP 0318216	A	31-05-1989	AU 2796789 A CN 1035679 A CN 1073719 A DD 287104 A DD 298524 A DD 298525 A DD 298526 A DD 298527 A DE 3886363 D DE 3886363 T DK 353789 A ES 2012739 T FI 893447 A FI 981380 A GB 2212511 A, B GR 90300069 T HK 38293 A HR 940493 A HU 9500427 A IE 62868 B IN 171237 A IN 171238 A IN 171239 A IN 171240 A IN 169067 A JP 2810032 B JP 10108674 A JP 10290696 A JP 10290697 A JP 2662358 B JP 6225797 A JP 2810022 B JP 9184844 A JP 9173079 A JP 2809956 B JP 6128290 A JP 2662350 B JP 6128291 A JP 2532805 B JP 6128292 A JP 2662351 B JP 6128293 A JP 2500880 T LV 10726 A LV 10726 B NO 955101 A NO 955102 A PT 89041 A, B SI 8812138 A US 5714596 A	14-06-1989 20-09-1989 30-06-1993 14-02-1991 27-02-1992 27-02-1992 27-02-1992 27-02-1992 27-01-1994 19-05-1994 18-07-1989 01-12-1994 17-07-1989 15-06-1998 26-07-1989 31-07-1991 30-04-1993 31-08-1997 28-12-1995 08-03-1995 22-08-1992 22-08-1992 22-08-1992 22-08-1992 31-08-1991 15-10-1998 28-04-1998 04-11-1998 04-11-1998 08-10-1997 16-08-1994 15-10-1998 15-07-1997 08-07-1997 15-10-1998 10-05-1994 08-10-1997 10-05-1994 11-09-1996 10-05-1994 08-10-1997 10-05-1994 29-03-1990 20-06-1995 20-12-1995 15-12-1995 15-12-1995 01-12-1988 31-08-1997 03-02-1998
EP 0388232	A	19-09-1990	AU 4750493 A AU 666767 B AU 666576 B	20-01-1994 22-02-1996 15-02-1996



# INTERNATIONAL SEARCH REPORT

Information on patent family members

Patent Application No

PCT/IB 98/01628

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0388232 A		AU 640920 B	09-09-1993
		AU 5278390 A	22-10-1990
		CA 2012482 A	17-09-1990
		DD 297446 A	09-01-1992
		FI 981381 A	15-06-1998
		GR 91300016 T	15-11-1991
		HU 9500746 A	29-01-1996
		JP 2693908 B	24-12-1997
		JP 7101986 A	18-04-1995
		JP 2809987 B	15-10-1998
		JP 7145194 A	06-06-1995
		JP 10309197 A	24-11-1998
		JP 10295389 A	10-11-1998
		JP 9215497 A	19-08-1997
		JP 2656995 B	24-09-1997
		JP 4504715 T	20-08-1992
		NO 303941 B	28-09-1998
		NO 960741 A	23-02-1996
		NO 960742 A	23-02-1996
		PT 93480 A,B	07-11-1990
		US 5714596 A	03-02-1998
		WO 9011089 A	04-10-1990
		US 5679342 A	21-10-1997
		US 5350671 A	27-09-1994
		US 5698390 A	16-12-1997
		US 5712088 A	27-01-1998
		US 5863719 A	26-01-1999
		US 5683864 A	04-11-1997
WO 9605513 A	22-02-1996	AU 3189495 A	07-03-1996
		CA 2174212 A	22-02-1996
		EP 0723665 A	31-07-1996
		JP 9504377 T	28-04-1997

# PATENT COOPERATION TREATY

From the:  
INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

To:

HALLYBONE, Huw George  
CARPMAELS & RANSFORD  
43 Bloomsbury Square  
London WC1A 2RA  
GRANDE BRETAGNE

## PCT

### WRITTEN OPINION

(PCT Rule 66)

Date of mailing  
(day/month/year)

**1 5. 09. 99**

Applicant's or agent's file reference

**P18098WO**

**REPLY DUE**

**within 2 month(s)**  
from the above date of mailing

International application No.

**PCT/IB98/01628**

International filing date (day/month/year)

**06/10/1998**

Priority date (day/month/year)

**06/10/1997**

International Patent Classification (IPC) or both national classification and IPC

**C12N15/12**

Applicant

**CHIRON S.P.A. et al.**

1. This written opinion is the first drawn up by this International Preliminary Examining Authority.

2. This opinion contains indications relating to the following items:

- I ☒ Basis of the opinion
- II ☒ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Rule 66.2(a)(ii) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain document cited
- VII ☒ Certain defects in the international application
- VIII ☒ Certain observations on the international application

3. The applicant is hereby invited to reply to this opinion.

**When?** See the time limit indicated above. The applicant may, before the expiration of that time limit, request this Authority to grant an extension, see Rule 66.2(d).

**How?** By submitting a written reply, accompanied, where appropriate, by amendments, according to Rule 66.3. For the form and the language of the amendments, see Rules 66.8 and 66.9.

**Also:** For an additional opportunity to submit amendments, see Rule 66.4.  
For the examiner's obligation to consider amendments and/or arguments, see Rule 66.4 bis.  
For an informal communication with the examiner, see Rule 66.6.

**If no reply is filed,** the international preliminary examination report will be established on the basis of this opinion.

4. The final date by which the international preliminary examination report must be established according to Rule 69.2 is: **06/02/2000.**

Name and mailing address of the international preliminary examining authority:



European Patent Office  
D-80298 Munich  
Tel. +49 89 2399 - 0 Tx: 523656 epmu d  
Fax: +49 89 2399 - 4465

Authorized officer / Examiner

**Julia, P**

Formalities officer (incl. extension of time limits)

**Vullo, C**

Telephone No. +49 89 2399 8061



**I. Basis of the opinion**

1. This opinion has been drawn on the basis of (*substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this opinion as "originally filed".*);

**Description, pages:**

1-28 as originally filed

**Claims, No.:**

1-26 as originally filed

**Drawings, sheets:**

1/17-17/17 as originally filed

2. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
- ☐ the claims, Nos.:
- ☐ the drawings, sheets:

3. This opinion has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

4. Additional observations, if necessary:

**see separate sheet**

**II. Priority**

1. ☐ This opinion has been established as if no priority had been claimed due to the failure to furnish within the prescribed time limit the requested:
- ☐ copy of the earlier application whose priority has been claimed.
  - ☐ translation of the earlier application whose priority has been claimed.
2. ☐ This opinion has been established as if no priority had been claimed due to the fact that the priority claim has been found invalid.

Thus for the purposes of this opinion, the international filing date indicated above is considered to be the relevant date.

**3. Additional observations, if necessary:****see separate sheet****V. Reasoned statement under Rule 66.2(a)(ii) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement****1. Statement**

Novelty (N)	Claims	1-18, 22-26; NO
Inventive step (IS)	Claims	1-26; NO
Industrial applicability (IA)	Claims	7

**2. Citations and explanations****see separate sheet****VII. Certain defects in the international application**

The following defects in the form or contents of the international application have been noted:

**see separate sheet****VIII. Certain observations on the international application**

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:

**see separate sheet**

**1. Additional remarks to item I :**

A "Sequence Listing" has been filed by the Applicant on 28.02.99. This "Sequence Listing" comprises SEQ ID No.: 1 to SEQ ID No.: 15 (pages 1-6).

**2. Additional remarks to item II :**

The priority documents pertaining to the present application were not available at the time of establishing this preliminary opinion. Hence, the current assessment is based on the assumption that all claims enjoy priority rights from the filing date of the priority document (06.10.97).

**3. Additional remarks to item V :**

The present application discloses that the DNA encoding the hepatitis C virus (HCV) receptor actually encodes the cellular protein known as CD81 (figure 1). This CD81 protein had been previously identified by monoclonal antibodies as the target of an antiproliferative antibody (TAPA-1) which inhibited in vitro cellular proliferation. The "EC2" region of CD81 (extracellular loop 2, second extracellular region of CD81) (residues 113-201) is the region involved in binding to the E2 protein of HCV. The application discloses the CD81 protein of chimpanzee, green monkey, hamster, rat and mouse too (figure 1). The application explicitly claims compounds that bind specifically to a CD81 protein (for use in the therapy or diagnosis of HCV), methods for screening such compounds, methods of treating, pharmaceutical compositions, uses in the manufacture of medicaments, assays for detecting HCV antibodies or HCV in a serum sample, diagnostic kits, transgenic non-human mammals, etc...

The following documents have been cited in the International Search Report (ISR) as being relevant for assessing the novelty and inventiveness of the claimed subject matter:

i) R. Oren et al. , Mol. Cell. Biol. 1990, Vol. 10 (8), pages 4007-4015 (D1) discloses the identification, cloning and characterization of the human 26-kd cell surface protein TAPA-1 (CD81) using a murine (cell) antiproliferative monoclonal antibody. The cDNA sequence and the corresponding deduced amino acid sequence are shown in figure 3. Reference is also made to an hybridization probe (coding and 3' untranslated region) used in genomic Southern blots for probing enzymatically digested genomic DNA. The deduced TAPA-1 amino acid sequence is said to be highly homologous to the leukocyte membrane protein CD37 and to the melanoma-associated antigen ME491. There is, however, no reference

in D1 to the possible use of the TAPA-1 antigen (CD81 protein) in the therapy or diagnosis of HCV.

ii) a similar disclosure is found in document M.L. Andria et al., J. Immunol. 1991, Vol. 147 (3), pages 1030-1036 (D2) which refers to the genomic organization and chromosomal localization of the TAPA-1 gene. D2 also refers to the homology of TAPA-1 to additional proteins, namely a cell surface marker on both haematopoietic and non-haematopoietic tissues (CD9), a pan-leukocyte antigen (CD53), a colon cancer antigen (CO-029) and a schistosome surface antigen (Sm23). Figure 3 outlines the relevance of the second extracellular loop of TAPA-1. There is, however, no reference in D2 to the possible use of the TAPA-1 antigen (CD81 protein) in the therapy or diagnosis of HCV.

In view of D1 and D2, the IPEA is of the opinion that the particular intended use disclosed in the application (i.e. therapy or diagnosis of HCV) must be seen as a distinctive characteristic of the claimed product which can not be disregarded for assessing the novelty and inventiveness of said claimed product (PCT Gazette-Section IV, Special Issue 29.10.98; PCT International Preliminary Examination Guidelines, Chapter III-4.8 and in particular Chapter IV-7.6) (first medical application). However, the IPEA is of the opinion that:

a) claim 4 is directed to a protein **comprising** the second extracellular region of CD81, i.e. the claim is not limited to this second extracellular region but it embraces any protein comprising said region and it is further not restricted to any intended use (therapy or diagnosis of HCV). The IPEA is of the opinion that the known TAPA-1 protein (CD81) certainly comprises this second extracellular loop and thus, D1 and D2 clearly anticipate the subject matter of claim 4 (Articles 33 (2) and (3) PCT).

b) claims 22-24 to general nucleic acid molecules hybridizing with a nucleic acid molecule which encodes a CD81 protein. In view of the nucleotide sequences and specific probes disclosed in D1 and D2, the IPEA is of the opinion that this subject matter is clearly anticipated by those documents (Articles 33 (2) and (3) PCT).

c) claim 8, 9 and 11 are directed to a general pharmaceutical composition comprising a CD81 protein or a protein comprising the second extracellular region of the CD81 protein and to a method for preparing said pharmaceutical compositions (no reference to HCV).

Claims 16-17 are directed to a general diagnostic kit comprising a CD81 protein but without any reference to HCV. In view of the high homology to well-known proteins and/or antigens which had already been used in general pharmaceutical compositions, diagnostic kits, etc... the IPEA is of the opinion that the skilled person would certainly have thought in using the TAPA-1 proteins in similar compositions. Thus, these claims do not seem to fulfil the requirements of Article 33 (3) PCT.

d) claims 19 and 20 are directed to a transgenic non-human mammal carrying a transgene encoding CD81 protein and a process for producing such a transgenic animal. The IPEA is of the opinion that once the nucleotide sequence of TAPA-1 was known and available, the production of a transgenic animal (mouse) would have not required any inventive contribution from the skilled person (Article 33 (3) PCT).

iii) WO-A-97/09349 (D3) discloses a human transmembrane cellular protein (putative cellular receptor) of 24-kD which is capable of specifically binding the E2 envelope protein of HCV. D3 further refers to pharmaceutical compositions comprising said protein, uses thereof for HCV diagnosis and therapy, methods for preparing such protein and for treating HCV, assay for detecting HCV antibodies, diagnostic kits, transgenic animals, etc... However, D3 actually discloses only the isolation of the 24-kD protein and its partial (functional) characterization but there is no amino acid sequence let alone a nucleotide sequence in D3.

In view of the (functional) characterization disclosed in D3, the IPEA is of the opinion that the protein isolated in D3 is the CD81 protein and that the actual contribution of the present application is the determination of its amino acid sequence and the nucleotide sequence of the corresponding gene and/or cDNA. However, the IPEA is of the opinion that once a protein is known and available to the skilled person, the determination of an inherent feature such as the amino acid sequence does not change any property of the known product per se and thus, it can not render novel this product. In addition, a document disclosing a (chemical) compound and its isolation and/or manufacture makes normally available this compound to the public in all desired degrees of purity (unless exceptional situations wherein all conventional purification processes have failed, etc...). The attention of the Applicant is also drawn to the fact that irrespective whether the protein disclosed in D3 is actually the claimed CD81 protein or not, the protein disclosed in D3 can certainly

**WRITTEN OPINION  
SEPARATE SHEET**

---

International application No. PCT/IB98/01628

be considered a "functional equivalent" of the CD81 protein (as it binds to the E2 of HCV). Thus, claims directed to "a CD81 protein or a functional equivalent thereof", namely pharmaceutical compositions, methods and uses, etc... as well as claims directed to a compound that binds specifically to a CD81 protein, namely the subject matter of claims 1-18 and 25-26 do not seem to fulfil the requirements of Articles 33 (2) and (3) PCT.

Furthermore, once a protein has been identified, the determination of a nucleic acid coding said protein does not seem to require any specific inventive skill, i.e. a the priority date a skilled person could expect to perform the cloning and expression of a gene in a fairly straightforward manner, and if the cloning, although requiring much work, does not pose such problems as to prove that the expectation of success was ill-founded, then inventive step can not be acknowledged. Thus, the IPEA is of the opinion that the subject matter of claims 19-24 does not fulfil the requirements of Article 33 (3) PCT over the teachings of D3.

iv) the documents EP-A-0 318 216 (D4) and EP-A-0 388 232 (D5) concern the characterization of the HCV, wherein the nucleotide sequence and amino acid sequence of the envelope region of the HCV (comprising the E2 region) are explicitly disclosed. Similarly the document D. Rosa et al., Proc. Natl. Acad. Sci. USA 1996, Vol. 93, pages 1759-1763 (D6) outlines the importance of the E2 region of HCV for target cellular binding. In this respect, WO-A-96/05513 (D7) emphasizes too the importance of the E2 region of HCV and its use for the detecting and identifying the cellular receptor for HCV. Thus, these documents seem to anticipate at least the subject matter of claim 6 (Articles 33 (2) and (3) PCT).

The attention of the Applicant is also drawn to the subject matter of claim 7 which is directed to a method of treatment of the human and/or animal body and thus, it may be excluded from examination by Article 34(4)(a)(i) PCT in combination with Rule 67(iv) PCT too. Furthermore, for such a subject matter no unified criteria exist in PCT for the assessment whether it is industrially applicable or not. The patentability can also be dependent upon the formulation of the claims. The EPO, for example, does not recognize as industrially applicable the subject matter of claims to the use of a compound in medical treatment, but will allow, however, claims to a known compound for first use in medical treatment (such as present claims 1-3, 5-6, etc...) and the use of such a compound for the



manufacture of a medicament for a new medical treatment (claims 12-13).

**4. Additional remarks to item VII :**

According to the ISA the Accession number of the CD81 protein in EMBL/GENBANK database is wrong. Accession number M33690 corresponds to rat alpha 2U globulin promoter region, whereas M33680 actually corresponds to human CD81.

If the Applicant intends to amend such an Accession number, the IPEA would like to draw its attention to the fact that even if it is obvious that an error has occurred in the application as originally filed, this is not enough for the allowableness of the rectification or the correction. In fact said rectification must be "per se" evident too (PCT Gazette-Section IV, Special Issue 29.10.98, "PCT International Preliminary Examination Guidelines", Chapter VI-7.14). If the Applicant fails to show a basis in the originally filed documents, this amendment will be considered to go beyond the disclosure as originally filed and thus, it will not be accepted under Rule 70.2(c) PCT.

**5. Additional remarks to item VIII :**

The following objections are also raised under **Article 6 PCT** concerning the clarity of the claims :

i) according to Article 6 PCT in combination with Rule 6.3 PCT the claims shall define the matter for which protection is sought in terms of technical features. The IPEA is of the opinion that a peptide, polypeptide, protein, oligonucleotide, gene, etc.. being chemical products must be clearly and unambiguously characterized by their amino acid and/or nucleic acid sequences, i.e. by reference to their specific SEQ ID No or Figure(s) comprising said sequences. The characterization of a product only by the desired function or by an arbitrary abbreviation without any real technical meaning (CD81) does not seem to fulfill the requirements of said Article 6 PCT in combination with Rule 6.3 PCT. Thus, this objection seems to apply for the subject matter of claims 1, 6-8, 10-12 and 14-26.

ii) the IPEA is of the opinion that the general reference to a "functional equivalent thereof" is ambiguous as far as said function is not clearly defined in the wording of the claim. A protein can have many "functions" depending on its intended use, such as for instance as

a substrate and/or inhibitor of particular enzymes (protease, methylases, etc...), an immunogen, foodstuff additive, etc... Even if the claim refers to the particular "use in the therapy or diagnosis of HCV", the IPEA is of the opinion that such use per se does not clearly and unambiguously characterize the meaning of the wording "functional equivalent", in particular in view of the definition found in the description, namely "any fragment or assembly of fragments of the complete protein that binds to HCV" (page 3 lines 7-9) (under suitable conditions (pH, salt, temperature, etc...) any peptide can bind to another polypeptide and wherein said peptide certainly comprises an arbitrarily short fragment (one, two, three residues) of any other peptide too) (the description refers to an homology of only 25% too !!). Thus, the scope of claims 1-2, 4, 7-8, 11-12, 14-16, 18-19, 21 and 25-26 seems to be ambiguous.

iii) in this respect too, the application discloses the presence and the sequence of CD81 proteins in hamster, rat and mouse (figure 1). These CD81 sequences are closely related, and they have a high homology (greater than 80%, with fragments being 100% homologous) to the CD81 sequences of human and chimpanzee. In particular, the CD81 protein from hamster has only 10 exchanged in the EC2 region (89 residues), the rat CD81 14 exchanged residues and the mouse CD81 only 17 residues (the Applicant is reminded, however, that most of the claims are directed to "a CD81 protein" and not to the specific "human and/or chimpanzee CD81 E2, known to be responsible for binding the HCV). As the Applicant rightly emphasizes, the "HCV receptor, whilst ubiquitous in humans and found in chimpanzees, is absent in other mammals" (page 12 lines 21-22). Thus, the IPEA is of the opinion that (a) whereas the CD81 proteins of these animals and fragments thereof are embraced by the general wording "a CD81 protein" and they are certainly "functional equivalents" of a CD81 protein, these "CD81 proteins" do not seem to be suitable, however, for the intended use (therapy or diagnostic of HCV). Furthermore, (b) transgenic animals comprising such "CD81 proteins" do not seem to be useful as "animal models of HCV infection" (see also example 7 on page 24, wherein it is also explicitly acknowledged that "human, but not the mouse EC2 loop of CD81, bound to E2 ... and inhibited binding of E2 to human cells").

iv) whereas there is no clarity problems for a claim directed to "a method for screening chemical compounds for ability to bind to the region of HCV responsible for binding to a host cell" wherein said method uses the CD81 protein (claim 18), clarity problems arise for claims directed to such products "per se". Claim 6 is only characterized by the result to be

**WRITTEN OPINION  
SEPARATE SHEET**

International application No. PCT/IB98/01628

achieved, namely "capable of binding specifically to a CD81 protein". The claimed product is only characterized by the desired functional property (desiratum) but not by any actual technical feature (parameters, structure, sequence, etc...). As far as said products are not clearly and unambiguously characterized, the IPEA is of the opinion that such a claim could embrace well-known products which under certain suitable conditions (pH, salt, temperature, etc...) could "specifically bind" the CD81 protein (the claim is not even limited to a protein, peptide and/or polypeptide but embraces any possible "compound"). The same objection applies too to the subject matter of claims 7-8 and 10-12.

v) in this respect, the description refers to the "ideal" compound that binds to CD81 as the one which only interacts with CD81 in the presence of HCV, so that the usual function of CD81 is not compromised on healthy cells. This requirement is not present in the wording of any claim related to such products, thus it is not clear whether a "compound" as claimed could be actually used "in the therapy or diagnosis of HCV" (claims 6-7, 10 and 12). In addition, it is not clear whether such "ideal" compounds could be easily screened by the method of claim 18, which only requires the presence of the chemical compound to be screened and the CD81 protein but in the absence of HCV.

vi) contrary to the subject matter of claim 15 which is directed to "an assay for HCV", the subject matter of claim 14 is directed to an assay for general "HCV antibodies". However, the reference to "HCV protein" in claim 14 seems to be ambiguous. There are many "HCV proteins" (produced by proteolysis of the HCV polyprotein) and it is not clear from said general wording whether all of them could be successfully used in the claimed assay or alternatively all of them should be used in the claimed assay.

vii) the subject matter of claims 19 and 20 refers to general "non-human transgenic mammals". The IPEA is of the opinion that such a broad subject matter is not fully (technically) supported by the present description. Methods for producing mammalian animals are not general for all mammals and they depend on the specific mammal used. Several steps such as a suitable status (stage) of the egg for optimal introduction of the DNA construct(s), time of insertion of the egg into the oviduct or uterus, etc.. are critical for the success of the method. Furthermore, several mammalian species do not lend themselves applicable to the present technology used for transgenic mice (or sheep, rabbits, goats and cows), such as duckbill platypus, spiny anteaters (echidna, egg-laying mammals, etc..). The IPEA is of the opinion that the description is not enabling for whole

**WRITTEN OPINION  
SEPARATE SHEET**

---

International application No. PCT/IB98/01628

range claimed, i.e. does not allow the skilled person to produce the claimed product in or for the whole range claimed (all mammalian species). The specification only teaches the production of a single mammalian species, namely mouse. It would be highly unpredictable and would require undue experimentation to prepare a homologous population of all mammalian species. The preparation of one species does not seem to be sufficient evidence to support that it can be prepared from all mammalian species.

viii) the wording of claim 22 seems to be completely ambiguous. There is neither a functional requirement nor a minimum length required for the claimed nucleic acid molecule. Furthermore, the claim does not provide any clear definition or characterization of the cited "standard conditions", which thus are completely ambiguous. In fact, the scope of the claim seems to embrace any (short) arbitrary nucleic acid (by selecting suitable "standard conditions") without bearing any technical relationship to the disclosed CD81 protein). The first part of this objection (function and length) also applies for the subject matter of claim 23 (and dependent claim 24).